

mortality from co-happening general-logical issues and unnecessary self destruction danger. In BD, risks for logical issues like diabetes or metabolic condition, and cardiovascular issues, and related mortality costs are various occurrences over the ones for the general people or with various mental issues.

Acknowledgement

None

Conflict of Interest

None

References

1. Fountoulakis KN (2010) The emerging modern face of mood disorders: a didactic editorial with a detailed presentation of data and definitions. *Ann Gen Psychiatry* 9: 14.
2. Sienaert P, Lambrichts L, Dols A, Fruyt J (2013) Evidence-based treatment strategies for treatment-resistant bipolar depression: a systematic review. *Bipolar Disord* 15: 61-69.
3. The UK ECT Review Group (2003) E f cacy and safety of electroconvulsive therapy in depressive disorders: a systematic review and meta-analysis. *The Lancet* 361:799-808.
4. Dierckx B, Heijnen WT, van den Broek WW, Birkenhager TK (2012) E f cacy of electroconvulsive therapy in bipolar versus unipolar major depression: a meta-analysis. *Bipolar Disord* 14:146-150.
5. Hiremani RM, Thirthalli J, Tharayil BS, Gangadhar BN (2008) Double-blind randomized controlled study comparing short-term e f cacy of bifrontal and bitemporal electroconvulsive therapy in acute mania. *Bipolar Disord* 10:701-707.