

Photodynamic Therapy with Verteporfin, a Novel Treatment Option for Retinal Arterial Macroaneurysm

V Swetha E Jeganathan^{1,2*} and Nitin Verma³

¹Department of Ophthalmology, The University Hospital Ayr, Scotland, UK

²The University of Edinburgh, Scotland, UK

³Hobart Eye Surgeons, Hobart, Tasmania, Australia

*Corresponding author: V Swetha E Jeganathan, Department of Ophthalmology, The University Hospital Ayr, Dalmellington Road, Ayr KA6 6DX, Scotland, UK, E-mail: vswetha@ausdoctors.net

Received date: Feb 26, 2016; Accepted date: May 13, 2016; Published date: May 20, 2016

Copyright: © 2016 Jeganathan VSE. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Abstract

Purpose: To report the novel use of photodynamic therapy (PDT) in a case of retinal arterial macroaneurysm (RAM).

Design: Interventional case report.

Methods: In the setting of institutional practice, a 76-year male patient presented with rapidly progressive vision loss in his left eye due to RAM. Three treatments of PDT were administered over 6 months.

Results: The patient's macroaneurysm regressed both clinically and angiographically during and following treatment with PDT.

Discussion: Our experience, albeit limited, nonetheless suggests that PDT may be a useful adjunct to current practice in the management of RAM, and certainly warrants more formal study.

Keywords: Photodynamic therapy; Retinal arterial macroaneurysm

Introduction

First described by Robertson in 1973 [1]. RAMs are acquired, localised dilatations of an arteriole within the radius of the 3rd branch of the retinal arterial tree. Current treatment options are controversial and of unproven benefit [2]. We present the case of a 76-year old gentleman with progressive vision loss from a unilateral RAM who responded favourably to treatment with PDT using Verteporfin. To the best of our knowledge, there are no previously published reports on the use of PDT in treating RAM.

Statement of Ethics

In accordance with the declaration of Helsinki, the patient provided informed consent prior to participation. We certify that all applicable institutional and governmental regulations concerning the use of human volunteers were followed during this research.

Material and Methods

A 76-year old, Caucasian gentleman with hypertension and type 2 diabetes mellitus noticed a sudden decline in his left vision. His visual acuity was 6/5 in the right eye and 6/12 in the affected eye. Slit lamp

Results

The patient received three treatments of PDT over 6 months. Over that time interval, his BCVA improved to 6/12 and the macroaneurysm regressed both clinically (Figure 2) and angiographically. His lesion most recently measured 2.488 mm² in diameter. No complications from PDT were noted.




Figure 2 Regressed macroaneurysm.

Discussion

Although the natural history of RAM is one of gradual and spontaneous involution with good visual prognosis in the majority of cases, a small percentage will progress to meaningful visual loss [4], secondary to haemorrhage, retinal oedema, or exudation involving the central macula. Released toxins, outer retinal shear forces, and a diffusion barrier created by hemorrhage are believed to contribute to photoreceptor damage and visual loss [5]. In our case, treatment became warranted when central vision loss occurred. Currently employed treatments for RAM are controversial and of unproven benefit [2]. Depending on the location of the RAM, these treatments can cause additional, significant functional damage. The use of PDT with Verteporfin for RAM has not been previously reported. Previous

studies have confirmed that PDT has been used effectively in treating age-related macular degeneration and several types of cancer and precancer. As the RAM involved the central macula, PDT was a safer option as it allows a selective occlusion of vascular lesions without damaging adjacent retinal structures. Although the exact mechanism of PDT is unknown, vascular-targeting PDT is believed to permeabilize blood vessels through the formation of endothelial intercellular gaps, which are postulated induced via endothelial cell microtubule depolymerization following vascular photosensitization