


Pre-Clinical Study: Regenerating autologous elastic cartilage for treatments of facial deformities using monkey auricular cartilage progenitor cells

Shintaro Kagimoto, MD PhD, Yokohama-city University Hospital, Yokohama, Japan

Abstract:

Statement of the Problem: Regenerative cartilage has a great potential to solve the present problem of facial surgery though there are several problems such as safety/quality/shape/volume of cartilage and cost. We recently identified a promising regenerative method based on the xenotransplantation of human auricular cartilage progenitor cells.



LWRU FHOOV WR LPPXQRGHILFLITQW DQLEPDOV .KEDIDVKE 7DNHEH HW DO 31\$6 DQG WKH DX Was 3DUidQives On Daye Dhas Problem s a Pa Next step.

key auricular cartilage progenitor cells to immunocompetent individuals (Kagimoto et al, Cell Transplant. 2016)

Biography:

to reconstruct self-renewing elastic cartilage, which may potentially be applied to patients with facial injuries and abnormalities. Today I will introduce our autotransplantation regenerative approach and the results. Methodology & Theoretical Orientation: We used 15 monkeys (Macaca fascicularis) and regenerated cartilage from their ear cartilage progenitor cells and autotransplanted them into their facial subcutaneous or submucosal regions to assume treatment surgery of present facial deformities.

Shintaro Kagimoto MD, PhD, a Japanese certified plastic surgeon and a certified regenerative medicine doctor at Yokohama-city university hospital, Japan. His speciality as clinical work is computer aided facial reconstruction for patients who have congenital or acquired facial deformity. He also conducts cartilage regenerative medicine / lymphedema research as-a medical researcher. He got a PhD there about cartilage tissue engineering using monkey ear cartilage progenitor cell (Kagimoto, 2016). He gained experience at Swansea Uni

We monitored the transferred tissue using non-invasive magnetic resonance imaging (MRI), assessed it histologically, and measured the glycosaminoglycan (GAG) levels. The experimental animal protocols were approved by the Animal Welfare and Animal Care Committee of the National Institute of Biomedical Innovation (Osaka, Japan) (approval ID: DS25-32). Findings: No complications such as inflammation or tumorigenesis were observed in all cases. The shape and volume of the transferred tissue remained, and also detected by MRI images. The harvested tissue were histologically assessed as elastic cartilage as well as their primary auricular cartilage and GAG levels were close to primary one. Conclusion & Significance: These approach could regenerate self-cartilage safely and promisingly. However, it still has difficulty to control the shape and volume of regenerative tissue in our-procedure. It might be necessary to add other methods such

Webinar On Plastic Surgery | September 10, 2020 | London, UK

Citation: Shintaro Kagimoto; Pre-Clinical Study: Regenerating autologous elastic cartilage for treatments of facial deformities using monkey auricular cartilage progenitor cells; Plastic Surgery 2020; September 10, 2020; London, UK