

Preoperative Indicators of Fruitful Weight Reduction Results Following Laparoscopic Sleeve Gastrectomy

Elizabeth Jung*

Department of Obstetrics and Gynaecology, Canada

Abstract

This study investigates preoperative indicators predictive of successful weight reduction outcomes in patients undergoing laparoscopic sleeve gastrectomy (LSG). As obesity rates continue to rise, bariatric procedures like LSG have become crucial interventions for weight management. Understanding the factors that contribute to favorable weight loss results is imperative for optimizing patient selection and counseling. A comprehensive analysis of preoperative markers, including demographic factors, comorbidities, psychological variables, and metabolic parameters, was conducted. The study employed a retrospective cohort design, analyzing data from a diverse population of patients who underwent LSG. Results reveal specific preoperative indicators that significantly correlate with successful weight reduction, providing valuable insights for clinicians in tailoring treatment plans and setting realistic expectations for patients undergoing LSG. The identification of predictive factors contributes to the enhancement of patient outcomes and the overall effectiveness of LSG as a bariatric surgical option.

*Corresponding author: Elizabeth Jung, Department of Obstetrics and Gynaecology, Canada, E-mail: ej.elizabeth@jung.com

Received: 03-Oct-2023, Manuscript No. jomb-23-117270; **Editor assigned:** 05-Oct-2023, PreQC No. jomb-23-117270 (PQ); **Reviewed:** 19-Oct-2023, QC No. jomb-23-117270, **Revised:** 25-Oct-2023, Manuscript No. jomb-23-117270 (R); **Published:** 31-Oct-2023, DOI: 10.4172/jomb.1000177

Citation: Jung E

Obtained ethical approval from the institutional review board (IRB) to ensure compliance with ethical standards for research involving human subjects. Ensured patient confidentiality and data anonymization during analysis. Extracted relevant data from electronic medical records, surgical databases, and patient charts. Employed a standardized data extraction form to maintain consistency

2. Horowitz M, Wilder S, Horowitz Z, Reiner O, Gelbart T, et al. (1989) The human glucocerebrosidase gene and pseudogene: structure and evolution. *Genomics* 4: 87-96.
3. Winfeld SL, Tayebi N, Martin BM, Ginns EI, Sidransky E, et al. (1997) Identification of three additional genes contiguous to the glucocerebrosidase locus on chromosome 1q21: implications for Gaucher disease. *Genome Res* 7: 1020-1026.
4. LeeCL, LeeKS, ChuangCK, SuCH, ChiuHC, et al. (2021) Otorhinolaryngological Management in Taiwanese Patients with Mucopolysaccharidoses. *Int J Med Sci* 18: 3373-3379.
5. Jilwan MN (2020) Imaging features of mucopolysaccharidoses in the head and neck. *Int J Pediatr Otorhinolaryngol* 134: 110022.
6. Murgasova L, Jurovcik M, Jesina P, Malinova V, Bloomfeld M, et al. (2020) Otolaryngological manifestations in 61 patients with mucopolysaccharidosis. *Int J Pediatr Otorhinolaryngol* 135: 110-137.
7. MacArthur CJ, Gliklich R, McGill TJI, Atayde AP (1993) Sinus complications in mucopolysaccharidosis IH/S (Hurler-Scheie syndrome). *Int J Pediatr Otorhinolaryngol* 26: 79-87.
8. Grabowski GA (2012) Gaucher disease and other storage disorders. *Hematology Am Soc Hematol Educ Program* 2012: 13-8.
9. Murugesan V, Chuang WL, Liu J, Lischuk A, Kacena K, et al. (2016) Glucosylsphingosine is a key biomarker of Gaucher disease. *Am J Hematol* 11: 1082-1089.
10. Bultron G, Kacena K, Pearson D, Boxer M, Yang M, et al. (2010) The risk of Parkinson's disease in type 1 Gaucher disease. *J Inherit Metab Dis* 33: 167-173.
11. Koprivica V, Stone DL, Park JK, Callahan M, Frisch A, et al. (2000) Analysis and classification of 304 mutant alleles in patients with type 1 and type 3 Gaucher disease. *Am J Hum Genet* 66: 1777-1786.