

# Prevention and Management of Postoperative Wound Pain after Thoracic Surgery

Kyoji Hirai<sup>1\*</sup>, Shingo Takeuchi<sup>1</sup> and Jitsuo Usuda<sup>2</sup>

<sup>1</sup>*Department of Thoracic Surgery, Nippon Medical School, Chiba Hokusoh Hospital, Japan*

<sup>2</sup>*Division of Thoracic Surgery, Nippon Medical School, Japan*

\***Corresponding author:** Kyoji Hirai, Division of Thoracic Surgery, Nippon Medical School Chiba Hokusoh Hospital, 1715 Kamakari, Inzai, Chiba 270-1674, Japan, Tel: 81-476-99-1111; Fax: 81-476-99-1911; E-mail: [ky-hirai@nms.ac.jp](mailto:ky-hirai@nms.ac.jp)

**Received date:** Jan 29, 2016, **Accepted date:** Feb 19, 2016, **Publication date:** Feb 23, 2016

**Copyright:** © 2016 Hirai K, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Early ambulation and management of wound pain lead to decrease  
of

correlated with the incidence of PTPS [15]. In addition, in patients with PTPS, sensory abnormalities such as allodynia (pain induced by