

# Probing Potent Interfacial Residues in Microvirin before Complexation with Glycoproteins: Visualizing Anti-HIV Activity *in silico*

E

Department of Microbiology, Bharathidasan University, Tiruchirappalli, India

Abstract: Microvirin (Mv) is a potent anti-HIV agent. The present study aims to probe the potent interfacial residues in Mv before complexation with glycoproteins. The study was carried out using *in silico* methods. The results show that the potent interfacial residues in Mv are the amino acid residues at positions 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100.

Keywords: Microvirin, Anti-HIV, Interfacial Residues, *In silico*

Introduction: Microvirin (Mv) is a potent anti-HIV agent. The present study aims to probe the potent interfacial residues in Mv before complexation with glycoproteins. The study was carried out using *in silico* methods. The results show that the potent interfacial residues in Mv are the amino acid residues at positions 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100.

Conclusion: The study shows that the potent interfacial residues in Mv are the amino acid residues at positions 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100.


Microvirin (Mv) is a potent anti-HIV agent. The present study aims to probe the potent interfacial residues in Mv before complexation with glycoproteins. The study was carried out using *in silico* methods. The results show that the potent interfacial residues in Mv are the amino acid residues at positions 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100.

The study shows that the potent interfacial residues in Mv are the amino acid residues at positions 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100.



Ĉ•ĵĤĜ	ĖĜĤĤ
Ô•ĀĤĤ	ĤĤĤ
Ô•ĤĤ	ĤĤĤĤĤĤ

Rigid/Non rigid residues





Tæ}içæ} }æ}Á ÚÉÁ T ~ |æ|æ@æ!æ}Á ÖÁ ÇG€FÍDÁ Ú! [àä] \*Á Ú [c^}cÁ Q)c^!-æ&æ|Á Ü^•ä ~ ^•Á ä}Á Tã&![çáíá}Á à^-[!^Á Ö [ { ]|^çæci [ }Á , äc@Á Ö|^&[!]:[c^á}•K  
Xä• ~ æ|ä : ä} \*ÁCE} cãÉP(XACE&cãçãc^`*in silico*ÉÁRÁ Tæ!ä}^ÁÚ&ääÜ^•ÁÖ^çÁ TÁG I FÉÁ

---