Biochemistry & Physiology: Open Access

Oladipo and Adebowale, Biochem Physiol 2017, 6:2
DOI: 10.4172/2168-9652.1000218

Research Article OMICS international

1SPEVDUJPO BOE 1BSUJBM 1VSJGJDBTUQJPSOHEM
OJHFSTPDJBUFE XJUI *MBKF -BLF 0OEP 4UBUF

Oladipo OO* and Olajide Adebowale

Department of Microbiology, Federal University of Technology, PMB 704, Akure, Nigeria

Abstract

7KH SUHVHQW VWXG\ DLPHG DW WKH LVRODWLRQ VFUHHQLQJ DQG SDUWLDO SX from soil and water samples collected from Ilaje Lake, Ondo state, Nigeria. The associated fungi were isolated DQG FRXQWHG E\ VWDQGDUG PLFURELRORJLFDO PHWKRGV 3DUWLDO SXUL¿FDWL

KLJKHVW VSHFL¿F PDQQDQDVH DFWLYLW\ ZDV UHFRUGHG IRU LVRODWH % ZKLOH 3XUL¿FDWLRQ RI FUXGH A. niger PDQQDQDVH ZDV DFKLHYHG RXW E\ DPPRQLXP VXOSKDWH SUHFLSLW (Sephadex G-200). Fractionation of ammonium sulphate precipitated mannanase from A. niger on Sephadex G-200 produced two activity peaks. In this investigation, fungal isolates evaluated for mannanase production from this source gave appreciable mannanase activity and this could be applied in many industrial processes.

Keywords: Beta-mannanase; Ilaje Lake; Locust bean gum; Fungal counts; Partial puri cation

Introduction

Beta-mannanase otherwise known as mannan endo-1,4- -mannosidase or 1,4- -D-mannanase with an enzyme commission (EC) 3.2.1.78 breakdowns -1,4-mannosidic bonds in the main chain of glucomannans, galactomannans and -1,4-mannans [1]. It transforms the abundant mannan-rich heteromannan, glucomannan and galactomannan substrates into manno-oligosaccharides [2] and a small amount of mannose, glucose and galactose [1,3]. A number of plants, bacteria, fungi and various invertebrates had been reported to produce -mannanase [1,3]. e mannanase of biological origins have found a variety of applications in di erent industrial sectors [3] including animal feed formulation, pharmaceutical products preparation, pulp biobleaching, [4] as well as pre-treatment of plant biomass for second generation biofuel production [5].

ere is a greater demand for biotechnological and industrial stable enzymes with higher catalytic e ciency that can function under harsh environmental conditions. In the recent years, screening and isolation of microorganisms for the production of industrial enzymes of microbial origin with unique and excellent characteristics has become one of the focuses of research in biotechnology. Enzymes required for industrial applications should be stable at high temperature, pH and presence of salts, solvents, etc. Extracellular enzymes from microbes associated with extreme environments such as halophiles are blessed with higher catalytic power to initiate and sustain metabolic activities under high salt concentrations. Gomes and Steiner [6] and Karan and Khare [7] have been reported that enzymes from extreme environments are expected to be active and stable under more than one extreme condition.

e physicochemical analysis carried out on four di erent coastal waters from Ilaje, Ondo state, Nigeria revealed that they were slightly acidic with a mean pH of 6.69 and also moderately saline with a mean salinity of 16.68% [8]. Hence, mannanase obtained from bacteria isolated from this source are expected to be halo tolerant. Mannanase producing microorganisms have been reported from di erent sources but there has been scanty or no information on mannanase production from bacteria inhabits Ilaje Lake. Hence, enzymes from halo tolerant bacteria from this location should have an edge in terms of stability. In search of high performance mannolytic organisms; we isolated and screened di erent fungi from soil and water samples in Ilaje, Ondo State, Nigeria. erefore, the present study dealt with the isolation and screening of fungal isolates for beta-mannanase production and its partial puri cation from Aspergillus niger

Materials and Methods

Sources of sample

Water and soil samples were collected from Ilaje Lake, Ondo State, Nigeria in sterile bottles. e samples were transferred to the laboratory and were used as sources for the isolation of mannanase-producing fungi [9,10].

Isolation and identi cation of mannanase-producing fungi from soil and water samples

e total fungal counts from the samples were determined using the pour plate method on potato dextrose agar. e samples were serially diluted and 1ml of an appropriate dilution was used to inoculate the plate in duplicate. e plates were incubated at 28 ± 2°C for 72 h, a er which the total colony count was determined as previously described

Corresponding author: Oladipo OO, Department of Microbiology, Federal University of Technology, P.M.B 704, Akure, Nigeria, Tel: +2348068054636; E-mail: mroladit@gmail.com

Received May 17, 2017; Accepted June 12, 2017; Published June 19, 2017

Citation: 2ODGLSR 22 $GHERZDOH 2 3URGXFWLRQ D Beta-Mannanase from Aspergillus niger Associated with Ilaje Lake, Ondo State, Nigeria. Biochem Physiol 6: 218. doi: 10.4172/2168-9652.1000218

Copyright: © 2017 Oladipo OO, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

could be attributed to the ability of their genetic makeup to secrete active mannanase coupled with varied di usion rate [10]. All the tested fungal strains were able to produce extracellular mannanase in submerged state fermentation, although with di erences in the rate of enzyme production. ese di erences might be attributed to the source of isolation and slight variation in their genetic makeup [10].

e variation in protein content generated by each of the strains

5.