



Qualitative Analysis of Unexplored Narratives

Bronte Marblemaw*

providing a more nuanced analysis [8,9].

Community participation: Involving communities connected to the unexplored narratives in the research process is essential. Collaborative approaches ensure that the analysis respects the authenticity of the narratives and incorporates the perspectives of those directly involved [10].

Narrative ethnography: Employing narrative ethnography allows researchers to immerse themselves in the context of the narratives. This approach goes beyond textual analysis, capturing the lived experiences and meanings embedded in the narratives.

Conclusion

The qualitative analysis of unexplored narratives is a vital endeavor in the pursuit of a more comprehensive understanding of human experiences. By valuing and analyzing narratives that have been marginalized or overlooked, researchers contribute to a more inclusive, diverse, and equitable representation of the tapestry of human stories. Embracing unexplored narratives is not only an academic pursuit but also a commitment to social justice, cultural appreciation, and a more enriched collective consciousness.

Conflict of interest

None

References

1. Torres AG (2004) Current aspects of Shigella pathogenesis. Rev Latinoam Microbiol 46: 89-97.
2. Bhattacharya D, Bhattacharya H, Thamizhmani R, Sayi DS, Reesu R, et al. (2014) Shigellosis in Bay of Bengal Islands, India: Clinical and seasonal patterns, surveillance of antibiotic susceptibility patterns, and molecular characterization of multidrug-resistant Shigella strains isolated during a 6-year period from 2006 to 2011. Eur J Clin Microbiol Infect Dis; 33: 157-170.
3. Von-Seidlein L, Kim DR, Ali M, Lee HH, Wang X, Thiem VD, et al. (2006) A multicentre study of Shigella diarrhoea in six Asian countries: Disease burden, clinical manifestations, and microbiology. PLoS Med 3: e353.
4. Germani Y, Sansonetti PJ (2006) The genus Shigella. The prokaryotes In: Proteobacteria: Gamma Subclass Berlin: Springer 6: 99-122.
5. Jomezadeh N, Babamoradi S, Kalantar E, Javaherizadeh H (2014) Isolation and antibiotic susceptibility of Shigella species from stool samples among hospitalized children in Abadan, Iran. Gastroenterol Hepatol Bed Bench 7: 218.
6. Sangeetha A, Parija SC, Mandal J, Krishnamurthy S (2014) Clinical and . J Health Popul Nutr 32: 580.
7. Nikfar R, Shamsizadeh A, Darbor M, Khaghani S, Moghaddam M. (2017) A Study of prevalence of Shigella species and antimicrobial resistance patterns in paediatric medical center, Ahvaz, Iran. Iran J Microbiol 9: 277.
8. Kacmaz B, Unaldi O, Sultan N, Durmaz R (2014) and clonality of sporadic Shigella sonnei isolates in Ankara, Turkey. Braz J Microbiol 45: 845-849.
9. Zamanlou S, Ahangarzadeh Rezaee M, Aghazadeh M, Ghotaslou R, et al. (2018) cephalosporinase, quinolone resistance, and molecular typing of Shigella spp. Infect Dis 50: 616-624.
10. Varghese S, Aggarwal A (2011) Extended spectrum beta-lactamase production in Shigella isolates-A matter of concern. Indian J Med Microbiol 29: 76.