



evolution' was used to say that the species able to survive is not the strongest but the most responsive to change. In this line, the aim of many scientists in the close future would be not only to collect data on the negative impact of climate changes on living biota but mainly to study as counteract the sources of thermal pollution together with identifying the rate of tolerance and adaptation capability of species to new climatic conditions.

## References

1. Byrne M, Przeslawski R (2013) Multistressor impacts of warming and acidification of the ocean on marine invertebrates' life histories. *Integr Comp Biol* 53: 582-596
2. Global temperatures. Met Office, UK.
3. Hansen J, Ruedy R, Sato M, Lo K (2010) Global Surface Temperature Change. *Reviews of Geophysics* 48:4: RG4004.
4. Comiso JC, Hall DK (2014) Climate trends in the Arctic as observed from space. *WIREs Climate Change* 5: 389-409
5. The Consequences of Global Warming On Glaciers and Sea Levels. Natural Resources Defense Council.
6. Houghton RA (2005) Aboveground forest biomass and the global carbon balance. *Global change biology* 11: 945-958
7. Gibbin EM, Chakravarti LJ, Jarrold MD, Christen F, Turpin V, et al. (2017) Can multi-generational exposure to ocean warming and acidification lead to the adaptation of life history and physiology in a marine metazoan? *J Exp Biol* 220: 551-563
8. Airi V, Gizzi F, Falini G, Levy O, Dubinsky Z, et al. (2014) Reproductive efficiency of a Mediterranean endemic zooxanthellate coral decreases