

**Keywords:** Tissue culture; Single nucleotide polymorphisms; Somaclonal variance; Genetic variants

**Index**

It is well known that plants have the capacity to regenerate via in vitro tissue culture from totipotent, differentiated somatic cells. Somaclonal variation is the word used to describe a phenomena that has been linked to a number of genetic and epigenetic instabilities, some



occurring mutations. This may be because all higher eukaryotes under study use the same highly conserved mechanism of deamination of methylated cytosines [13].

When considered as a whole, it is evident that the types, rates, and spectra of tissue culture-induced mutagenesis are highly variable and may differ depending on the species, genotypes, and tissue culture conditions, or may simply be accidental. Both previous studies and our data here showed that at least TE mobilisation occurs concurrently with DNA methylation dynamics, consistent with their repressive control by this epigenetic marker. This is contrary to our previous finding based on DNA marker analysis that genetic mutation is the major type of molecular changes associated with rice tissue culture. Given that tissue culture induced mutagenesis is known to occur stochastically, the participation of epigenetic processes supports the fortunate component of this process [14].

## Conclusion

Prior research has discovered domestication-related rice areas that are genetically brittle and vulnerable to mutations in the presence of natural selection. Here, we demonstrated that there was minimal correlation between the genomic areas that were hyper-mutagenic in tissue culture and those that were hyper-mutable in the wild. This implies that natural mutation and tissue culture-induced mutagenesis may have different mutagenic pathways. Of fact, since the somaclonal line (TC-reg-2008) we utilised has only experienced eight generations without deliberate selection, we cannot completely rule out the notion that long-term selection was what made the difference. However, we believe that further research is necessary to elucidate any novel potential applications of tissue culture in crop development.

## Acknowledgement

None

## Conflicts of Interest

None

## References

- Krishna H, Sairam RK, Singh SK, Patel VB, Sharma RR, et al. (2008) Mango explant browning: Effect of ontogenic age, mycorrhization and pre-treatments. *Sci Hort* 118: 132-138.
- Uchendu EE, Paliyath G, Brown DC, Saxena PK (2011) In vitro propagation of North American ginseng (*Panax quinquefolius*, p. L.). *In Vitro Cellular & Developmental Biology-Plant* 47: 710-718.
- Laukkanen H, Rautiainen L, Taulavuori E, Hohtola A (2000) Changes in cellular structures and enzymatic activities during browning of Scots pine callus derived from mature buds. *Tree Physiol* 20: 467-475.
- Aliyu OM (2005) Application of tissue culture to cashew (*Anacardium occidentale*, p. L.) breeding: An appraisal. *Afr J Biotechnol* 4: 1485-1489.
- Panaia M, Senaratna T, Bunn E, Dixon KW, Sivasithamparam K (2000) Micropropagation of the critically endangered Western Australian species, *Symonanthus bancroftii* (F Muell.)L. Haegi (Solanaceae). *Plant Cell Tissue Organ Cult* 63: 23-29.
- Beckman CH (2000) Phenolic-storing cells: keys to programmed cell death and periderm formation in wilt disease resistance and in general defence responses in plants? *Physiol Mol Plant Pathol* 57: 101-110.
- Amin Dalal M, Sharma BB, Srinivasa Rao M (1992) Studies on stock plant treatment and initiation culture mode in control of oxidative browning in in vitro cultures of grapevine. *Sci Hort* 51: 35-41.
- Tang W, Harris LC, Outhavong V, Newton RJ (2004) Antioxidants enhance in vitro plant regeneration by inhibiting the accumulation of peroxidase in Virginia pine (*Pinus virginiana* Mill.). *Plant Cell Rep* 22: 871-877.
- Tang W, Newton RJ, Outhavong V (2004) Exogenously added polyamines recover browning tissues into normal callus cultures and improve plant regeneration in pine. *Physiol Plant* 122: 386-395.
- Thomas TD (2008) The role of activated charcoal in plant tissue culture. *Biotechnol Adv* 26: 618-631.
- Shukla MR, Jones AMP, Sullivan JA, Liu C, Gosling S, et al. (2012) In vitro conservation of American elm (*Ulmus americana*): potential role of auxin metabolism in sustained plant proliferation. *Can J Forest Res* 42: 686-697.
- Zo J, Amrhein N (1992) Inhibitors of Phenylalanine Ammonia-Lyase: 2-Aminoindan-2-phosphonic Acid and Related Compounds. *Liebigs Ann Chem*: 1992: 62-628.
- Grabber JH, Hatfeld RD, Ralph J, Zon J, Amrhein N (1995) Ferulate cross-linking in cell walls isolated from maize cell suspensions. *Phytochemistry* 40: 1077-1082.
- Mauch-Mani B, SlusarenS I-B0048004Cphosphoni CH (e: 10718indan. *Liebigs Ann*