

Short Note on Orthopedics

Kattekola Prasanna Laxmi*

Department of Pharmacology, Osmania University, Hyderabad, India

Abstract

Youthful Adults with hip torment whose radiographs are typical may have one of three snapping-hip conditions that can be treated with satisfying outcomes. These disorders ought to be separated from an easy profound "pop" that happens with typical hip movement and has no clinical significance. The snapping-hip disorders happen regularly in the age bunch 15 through 40 years and are marginally more normal in ladies.

Keywords:

Introduction

The hip joint is a ball-and-socket joint that allows for a wide range of motion. It is composed of the femoral head of the femur and the acetabulum of the pelvis. The hip joint is surrounded by a capsule and is supported by several ligaments. The hip joint is a weight-bearing joint and is subject to a great deal of stress and strain. Hip pain is a common complaint and can be caused by a variety of factors, including trauma, degenerative changes, and inflammation. Hip pain can be a significant source of disability and can affect a person's quality of life. The diagnosis of hip pain is often challenging and requires a thorough history and physical examination, as well as imaging studies such as X-rays, MRI, and CT scans. The treatment of hip pain depends on the underlying cause and may include conservative measures such as rest, physical therapy, and pain medications, as well as surgical options such as hip replacement surgery.

.....

Conclusion

.....