



Membrane-anchored NRG-1 is cleaved by presenilin-1

Membrane-anchored NRG occurs abundantly in the prefrontal cortex, hippocampus, cerebellum and substantia nigra [23-28]. NRGs are expressed by various immune cells, such as astrocytes, oligodendroglial cells, microglial cells, and neurons, in the brain [25,29-31]. NRGs are cleaved by proteases into presenilin-1 or

are

-
28. Thuret S (2004) The neuregulin receptor, ErbB4, is not required for normal development and adult maintenance of the substantia nigra pars compacta. *J Neurochem*: 1302-1311.
 29. Linying, Z (2014) Neuroprotective effects of neuregulin-1 ss on oligodendrocyte type 2 astrocyte progenitors following oxygen and glucose deprivation. *Pediatr Neurol* 50: 357-362.
 30. Wen L (2010) Neuregulin 1 regulates pyramidal neuron activity via ErbB4 in parvalbumin-positive interneurons. *Proc Natl Acad Sci* 107: 1211-1216.
 31. Jaworski A, Burden SJ (2006) Neuromuscular synapse formation in mice lacking motor neuron- and skeletal muscle-derived Neuregulin-1. *J Neurosci* 26: 655-661.
 32. Mei L, Xiong WC (2008) Neuregulin 1 in neural development, synaptic plasticity and schizophrenia. *Nat Rev Neurosci* 9: 437-4352.
 33. Gambarotta G (2004) ErbB4 expression in neural progenitor cells (ST14A) is necessary to mediate neuregulin-1 beta1-induced migration. *J Biol Chem* 279: 48808-48816.
 34. Pitcher GM (2011) Schizophrenia susceptibility pathway neuregulin 1-ErbB4 suppresses Src upregulation of NMDA receptors. *Nat Med* 17: 470-478.
 35. Hahn CG (2006) Altered neuregulin 1-erbB4 signaling contributes to NMDA receptor hypofunction in schizophrenia. *Nat Med* 12: 824-828.
 36. Neddens J (2009) Neuregulin links dopaminergic and glutamatergic