

## Abstract

Wheat is one of the most important cereal crops in Ethiopia and produced across large area of the country. Production of the crop constrained by several infection diseases including rust and Septoria leaf blotch diseases which are the major bottle neck of wheat production in Ethiopia. The objective of this study was to assess distribution,

analysis was carried out by inoculating single uredinial isolates on to the 20 differential hosts. The result indicated that, stem, leaf and yellow rust mean incidence value 54.7%, 19.4% and 7.7% were recorded in the surveyed areas, respectively and mean severity value of 7.0%, 9.7% and 5.5% in the same order. Septoria leaf blotch was the most prevalent disease with 100%. Mean incidence of 83% and 0.44 disease index of Septoria leaf blotch were recorded. The most widely grown Varieties Dig TJ/TT1 1 Tf0.037 Tw T\*[95%. Stem rust resistance )-0.6(genes )]TJ/TT2 1 Tf0 Tw 15.137 0 Td(Sr)

---

\*Corresponding author: Endale Hailu, Ethiopian Institute of Agricultural Research, Ambo Plant Protection Research Center, P.O. Box 37, Ambo, West Shewa, Ethiopia, Tel: 25191363779; E-mail: [sebhailuabera@yahoo.com](mailto:sebhailuabera@yahoo.com)

Received March 10, 2015; Accepted April 02, 2015; Published April 04, 2015


Citation: Hailu E, Woldeab G (2015) Survey of Rust and Septoria Leaf Blotch Diseases of Wheat in Central Ethiopia and Virulence Diversity of Stem Rust *Puccinia graminis* f. sp. *tritici*. Adv Crop Sci Tech 3: 166. doi:[10.4172/2329-8863.1000166](https://doi.org/10.4172/2329-8863.1000166)

Copyright: © 2015 Hailu E, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.









### Acknowledgment

We would like to offer a great thanks to wheat rust research team of Ambo plant protection research Center for their valuable encouragement and technical support during the whole period of the study.

### References

1. Hailu G, Tanner DG, Mengistu H (2011) Wheat research in Ethiopia: A Historical perspective, IARI and CIMMYT, Addis Ababa.
2. Bekele, Verkuij H, MWangi W, Tanner D (2000) Adoption of improved wheat technologies in Adaba and Dodola woredas of the Bale high lands, Ethiopia, Mexico.
3. CSA (central statistics Authority) (2013) Report on area and crop production forecast for major grain crops, Addis Ababa, Ethiopia: statistical bulletin.
4. Zegeye T, Taye G, Tanner D, Verkuij H, Agidie A, et al. (2001) Adoption of improved bread wheat varieties and inorganic fertilizer by small-scale farmers in Ethiopia. CIMMYT, Mexico city, Mexico.
5. Prescott JM, Burnett PA, LeSaari EE, Ranson J, Bowman J, et al. (1986) Yemane, T. A review of research on diseases of berley, tef and wheat in Ethiopia. In: Tsedeke Abate (ed.), A review of crop protection research in Ethiopia. Institute of Agricultural Research (IAR), Ethiopia, pp 79-107.
6. Bekele E (1985) A review of research on diseases of berley, tef and wheat in Ethiopia. In: Tsedeke Abate (ed.), A review of crop protection research in Ethiopia. Institute of Agricultural Research (IAR), Ethiopia, pp 79-107.
- 7.