Susceptibility of Snails to Infection with Schistosomes is influenced by Temperature and Expression of Heat Shock Proteins

1,3* 1 1 1 2 2 1 1 1 3 3 2 3

[1]*University of the District of Columbia, 4200 Connecticut Ave, Washington, D.C. 20008, USA*

2

of susceptible NMRI snails with the Hsp 90 inhibitor, geldenamycin, rendered these snails resistant [18].

Heat shock proteins (Hsps) are evolutionarily highly conserved molecular chaperones that function to protect the cell during stressful conditions. Hsps are actively synthesized in response to any cellular stress from heat shock, infection, or trauma. Their ability to bind to mis-folded and newly synthesized proteins during stress prevents protein aggregation, and therefore, cell-death. Although Hsps are intracellular proteins, they also are released in exosomes but possibly

Containing a recognition site for BamH1, a 5'overhang to ensure proper cutting by the restriction enzyme and a spacer (T) to keep the ORF in frame and a 3' primer (hsp 70-2exp3'):

5'-CGG GAATTC TCA ATCTACTTCTTCAACTGTTGGGCC-3' (36-mer)

overhang EcoRI stop - HSP 70 3'ORF (24 bp)

Containing a recognition site for EcoR1, a 5'overhang to ensure proper cutting by the restriction enzyme and a stop codon. A three step-cycle PCR program of 94°C for 1 min, 65°C for 1 min and 72°C for 1.5 min was employed for 30 cycles. The gel-purified PCR product of 1930 bp was digested with BamH l and EcoR1 and ligated into the expression vector pRSET B (Invitrogen, Carlsbad, CA) in frame. The recombinant plasmid was used to transform *E. coli* (DH10B). In order to express the recombinant protein from the BgHsp 70-pRSETB constructs, purified recombinant DNA from the transformants propagated in *E. coli* DH10B were checked for the correct size and orientation (in-frame) and verified by PCR prior to transformation into *E. coli* B strain BL21 (DE3) pLYsS (Invitrogen, Carlsbad, CA). Protein expression in the transformants containing the BgHsp 70 insert was induced with 0.4 mM IPTG for 2 h at 37°C according to manufacturer's instructions (Invitrogen, Carlsbad, CA). The expressed BgHsp 70-His-tagged fusion protein was purified using a nickel-NT agarose 158 column. For elution, buffer contained varying concentrations of imidazole (60 mM-500 mM) according to the manufacturer's recommendation (Novagen/EMD Biosciences, Madison, WI). The recombinant protein, eluted in 200 mM imidazole, was dialyzed against Tris-buffered saline (TBS, pH 8.0), and quantitated by the bicinchoninic protein assay (Pierce, Rockford, IL). Antibodies were produced in six Swiss Webster mice (Charles River Laboratories Inc., Wilmington, MA) using purified recombinant BgHsp 70 protein. Approximately 50 µl of blood was obtained from

the individual snails when they were first introduced at the higher temperature were discarded. All parasite exposures of 32°C maintained BS-90 (BS- 90_32°C) was undertaken at 25°C. These schistosome-exposed BS-90_32°C progeny (F1 to F4) snails were either maintained at room temperature or returned after exposure to 32°C to monitor for shedding of cercariae. In addition, BS-90_32°C progeny after two hours' exposure were frozen and used for protein extraction. As shown in Figure 3A, all the 32°C maintained progeny BS-90 (100% from both experiments) even as early as the first generation (F1) or later (F4, data not shown) were found to be positive at four weeks post-exposure, within the same time frame as the highly susceptible NMRI snail. Intriguingly, all the BS-90_32°C progeny that were returned to 32°C after exposure failed to survive (data not shown). Although all the exposed BS-90_32progeny (F1 to F3) were found to be positive for infection as shown in Figure 3B, they however, shed fewer cercariae than exposed susceptible NMRI snails. Room temperature (25°C) - BS-90 snails that were exposed at the same time as the 32°C progeny snails failed to shed cercaria and, as anticipated, were resistant to *S. mansoni* (Figures 3A and 3B).

Figure 2



Figure 2: The percentage of BS-90 snails surviving after incubation over time (months) at 32°C. Note the rapid decline in mortality occurred within the first month of introducing the snails to 32°C.



Figure 3A: Histograms showing percentage of susceptible NMRI, BS-90_32 Progeny, F1, F2 and F3 shedding cercaria after 4 weeks post-exposure to *S. mansoni* miracidia. Note that the non-infected control (unexp) BS-90 room temperature (RT) resident snail, exposed (Exp) in parallel to the same number of miracidia, remained negative for infection.



Figure 3B: Histograms showing the number of cercariae shed from susceptible NMRI and BS-90_32 F1, F2 and F3 progeny snails. Note that the non-infected control (unexp) room temperature (RT) resident BS-90 exposed in parallel to the same number of miracidia remained negative for infection.

Susceptible and resistant juvenile snails display different levels of Hsp 70 in response to either heat shock or infection with *S. mansoni*

Resistant BS-90 and susceptible NMRI snails respond differently in the temporal kinetics and levels of Hsp 70 mRNA induced after either heat shock at 32°C or after early exposure to miracidia [21]. Here we quantified the Hsp 70 protein by ELISA in protein extracted at different time points after exposure to schistosome miracidia or heat shock for 15, 30, 120, or 240 minutes from NMRI susceptible (Figure 4A) and BS-90 resistant (Figure 4B) snails.

As shown in Figure 4 either heat shock or *S. mansoni* infection induced enhanced expression of Hsp 70 in the susceptible NMRI (Figure 4A) than the resistant BS-90 (Figure 4B) at all-time points. Compared to the susceptible snail where Hsp 70 protein expression was observed early after schistosome infection (30 min), Hsp 70 protein induction was low and slow (120 min) in the resistant snail. Interestingly, in the susceptible NMRI snail higher levels of Hsp 70 protein were detected following *S. mansoni* infection than after the heat shock. Analysis using ELISA of protein extracts from these groups of snails with the pre-immune antiserum, as control, detected no



Figure 4: A) Histograms showing absorbance at OD495 of susceptible NMRI snail soluble protein extracts in an ELISA assay using mouse antiserum raised against the purified snail recombinant Hsp 70 protein expressed in *E. coli.* Protein extracts were prepared from snails, either after heat shock at 32°C or following exposure for different time point p M



Figure 6: A) Western blot analysis of soluble protein extracts prepared from juvenile susceptible (NMRI) and resistant (BS-90) snails after either heat shock (32°C) or following exposure. for different time points to normal miracidia using anti-Hsp 70 serum as described in Materials and Methods. Note that there is significant expression of Hsp 70 at early time points after normal schistosome exposure in the susceptible NMRI compared to the expression of this protein in the resistant BS-90 snail. Arrows are used to indicate the lanes loaded with protein samples from heat shocked (Hsp) vs. schistosome exposed (Exp) snails. B) Western blot analysis of soluble protein extracts prepared from juvenile susceptible (NMRI) and resistant (BS-90) snails after either heat shock (32°C) or following exposure for different time points to radiation-attenuated miracidia using anti-Hsp 70 serum as described in Materials and Methods. Note expression of Hsp 70 after attenuated parasite exposure in both susceptible NMRI compared to the resistant BS-90 snail remains the same at all-time points examined.



Figure 6C: Western blot analysis of soluble protein extracts prepared from adult susceptible (NMRI) and resistant (BS-90) snails after either heat shock (32°C) or following exposure for different time points to normal miracidia using anti-Hsp 70 serum as described in Materials and Methods. Note expression of Hsp 70 after heat shock in susceptible adult NMRI is significant at all-time points when compared to expression of this protein in resistant adult BS-90 snails. Hsp 70 protein expression in adult BS-90 snails, post- exposure at all-time points was negligible. Arrows are used to indicate the lanes loaded with protein samples from heat shocked (Hsp) vs. schistosome exposed (Exp) snails.

Discussion

Previously we had reported that susceptibility of the intermediate snail host, *B. glabrata* to *S. mansoni* infection, is influenced by elevated temperature. Moreover, we also showed that an early induction of stress was a characteristic feature of susceptible but not resistant snails. Therefore, treatment of the susceptible NMRI snail with the Hsp 90 inhibitor, geldenamycin, rendered this snail non-susceptible to *S. mansoni* [18,28]. In this follow up study, we have shown that breeding the resistant BS-90 snail at 32°C rendered all progeny (F1 and F4) produced in this warmer environment susceptible to *S. mansoni* after exposure at 25°C. The BS-90 snail are normally resistant at ambient temperature and have been considered for the past two decades as the representative resistant snail strain that is used in most molecular studies aiming to identify genes that underscore the infection refractory phenotype [18].

The BS-90 stock is a wild type pigmented snail that was identified in Salvadore, Brazil by Correa and Paraense [29]. At room temperature (25°C), this snail is refractory at all ages (juvenile and adult) to both new and old world *S. mansoni*, reliably destroying the invading miracidia by an active innate defense mechanism where hemocytes and plasma factors quickly kill the larval schistosome [30-32].

For the susceptible NMRI stock used in this study, however, early recruitment of hemocytes that engulf the parasite fails to manifest and the miracidia establish the infection and develop into sporocysts stages that by asexual reproduction produce cercariae that are infectious to the human host. Understanding the complex mechanism(s) behind these diverse outcomes of the schistosome/snail host interaction (notwithstanding the parasites own genetic make-up that also shapes this relationship) has been the driving force for developing new vector-based strategies to control schistosomiasis. In this regard, the anticipation has been to utilize resistant snails either to displace susceptible ones in the field to block transmission of schistosomiasis, or to use resistant snails to elucidate molecular pathways that underlie refractoriness to the schistosome trematode in its intermediate snail host [15,16].

Investigation of differential gene expression between BS-90 and