Synthesis of Dihydroimidazole Derivatives under Solvent Free Condition and Their Antibacterial Evaluation

Mahmoud Abd El Aleem Ali Ali El-Remaily[1*], Shaaban K Mohamed[2], Ahmed MM Soliman[1], and Hossam Abd el-Ghanya[1]

[1]*Department of Chemistry, Sohag University, Egypt*

[2]*School of Chemistry and Environmental Science, Manchester Metropolitan University, Manchester, England*

[*]**Corresponding author:** Mahmoud Abd El Aleem Ali Ali El-Remaily, Department of Chemistry, Sohag University, Sohag 82524, Egypt, Tel: +20108036348; Fax: +2093601159; E-mail: msremaily@yahoo.com

Rec date: Jul 03, 2014, **Acc date:** Jul 25, 2014, **Pub date:** Aug 01, 2014

Copyright: © 2014 Ali El-Remaily MAEAA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Reaction of 2-guanidinobenzoxazole with several halogenated active methylene compounds has revealed formation of the corresponding dihydroimidazole derivatives under solvent free conditions in excellent yield. The anti-bacterial evaluation of the newly synthesized products against different types of Gram-positive and Gram-negative bacteria was performed. Most of products showed high inhibitory effect. The structure of all compounds has been characterized by on IR, 1H-NMR, 13C-NMR, Mass spectrum and some X-ray diffractions.

Keywords: Guanidylbenzoxazol; Dihydroimidazole, Halogenated active methylenes; Anti-bacterial

Introduction

Imidazole containing drugs exhibit different types of

was heated under reflux for 3 hours. The reaction mixture was allowed to cool down then treated with 10% NaOH to afford the pure product 1 in 98% yields.

General procedure for preparation of compounds 2 – 7

A mixture of compound 1 (50 mmol) and 10 ml of halogenated active methylene compounds such as phenacyl bromide, chloro acetyl chloride, chloro acetone, ethyl bromo acetate, chloro acetonitril and bromo malononitril, in addition to few drops of catalytic glacial acetic acid was refluxed for 4 hours. The resulting solid product was collected by filtration, dried under vacuum, washed with cold ethanol and recrystallized from ethanol in excellent yield.

Synthesis of N-[(2Z)-4-phenyl-1,3-dihydro-2H-imidazol-2-ylid-ene]-1,3-benzoxazol-2-amine 2

Yield 90%, mp 255°C ; IR: cm^{-1} 3288, 3178 (2NH); 1HNMR: δ 09.74 (