The Expression of S100A1 Protein and its Usage for Differential Diagnosis between Normal Ovarian Tissue and Ovarian Cancer

Angel Danchev Yordanov[1], Borislav Alioshev Ignatov[2], Borislava Ivova Dimitrova[1], Ivan Nedkov Ivanov[3], Stanislav Hristov Slavchev[4], Momchil Denev Ivanov[5] and Polina Petkova Vasileva[6] March 30, 2017; **Accepted** April 02, 2017; **Published** April 08, 2017

Citation: Yordanov AD, Ignatov BA, Dimitrova BI, Ivanov IG0M 8S-C1i0 1 T i4 i4 iaa et al.Yord The Expression of S100A1 Protein and its Usage for Differential Pub-24

100 1

Aim

100 1

.

' 0

(‘0 ‘0),

,

" . " - .

100 1 .

Introduction

100 21 .

100 1 , .

, , 1-5 .

, 1-10 .

100 1

. .

, 100 1 11

.

.

100 1 12 .

,

100 1.

. 100 1

, .

60% ,

(6) 100 1.

100 1

, ,

. 2 , 1 0.0 . 20 (64.52%) ,

(5 5%) .

1 2

(66.6 %). 1

(%) 100% (1 1)

(11 12).

11(%)

( ).

100 1

2 0.0 , 2 0. 0 (2).

Discussion

, 100 1

-

. , .

2(%) . 5 (15.62%)

.

(2)

, .



Figure 11: Coloring in serous papillary ovarian cancer.

Figure 12: Coloring in mucinous ovarian cancer.

	Low grade (G1)	Intermediate grade(G2)	High grade (G3)
Without S100A1expression	2 (5.00%)	4 (10.00%)	3 (7.50%)
With S100A1 expression	6 (15.00%)	14 (35.00%)	11 (27.50%)

Table 2: S100A1 expression in different tumor grades.

.

, , , ,

.

100 1

. (

)

.

- ,

, , .

.

, .

,

100 1. 100 1

(.6%) ,

(6 . %).

( )0.5 (00 0.2 0 -1.2 , 0.(( )0.6(02 0.0 5 (