

¹, Valentin Brodszky ², Zsombor Zrubka ², Attila Zaránd ³, Tüde Mezei ⁴, Zsuzsanna Szilasi ⁵, Adrienne Fehé ⁶, Iván Igaz ⁷, Krisztián Somlai ⁸, Keresztély Merkel ⁹ and Zsolt Baranyai ^{3*}

¹Department of Otorhinolaryngology and Head and Neck Surgery, Jahn Ferenc Hospital, Budapest

²Department of Health Economics, Corvinus University of Budapest, Hungary

³1st. Department of Surgery, Semmelweis University, Hungary

⁴Department of Urology, Jahn Ferenc Hospital, Budapest, Hungary

⁵Department of Otorhinolaryngology and Head and Neck Surgery, HDF Medical Centre, Budapest, Hungary

⁶Department of Laboratory Medicine, Semmelweis University, Hungary

⁷Department of Gastroenterology, Szent Imre Hospital, Hungary

⁸Department of Surgery, Szent Margit Hospital, Hungary

⁹Department of Surgery, Szent Imre Hospital, Budapest, Hungary

Abstract

Introduction: IL-6 is an essential factor in inflammatory processes; its level has been found to be elevated in a variety of gastrointestinal tumors. Nearly 30% of IL-6 circulating in the blood is produced by adipose tissue; however, there are no clear data on whether its level changes in obese patients.

Material and method: We studied 127 patients with gastrointestinal tumors of different stages. IL-6 levels, white blood cell count (WBC), and hemoglobin (Hgb) concentration were determined in preoperative blood samples. BMI was determined with a calibrated calculator, before surgery. During the statistical analysis of the data, correlation was calculated using Spearman, Wilcoxon, and Pearson correlation coefficients.

Results: Valéria Jósa

Keywords: BMI; Nutritional status; IL-6; Gastrointestinal tumor

Introduction

Neoplastic disease and obesity are among the most alarming public health problems of developed countries [1,2]. According to the literature, obesity and the increase of the body-mass index (BMI) are associated with an enhanced risk of esophageal, colorectal, pancreatic, and gallbladder neoplasms, as well as of postmenopausal breast tumors and endometrial cancer [3-8]. In overweight (BMI 25 to 29 kg/m²) and in obese (BMI ≥ 30 kg/m²) patients, the incidence of these tumors increases from 3% to 10% [9]. The exact underlying pathomechanism is unknown. Elevated serum IL-6 levels were observed in patients with various gastrointestinal tumors [10-12]. In the neoplastic microenvironment, the primary sources of IL-6 include tumor cells, tumor-associated macrophages (TAM), CD4⁺ T cells, myeloid-derived suppressor cells (MDSC), and fibroblasts [13-16]. IL-6 has a regulatory role in nearly every process of tumorigenesis: it inhibits apoptosis [17,18], as well as facilitates survival [19,20], proliferation [21,22], and angiogenesis [23]. Further, it enhances invasiveness and metastasis formation [16,24], and stimulates the metabolism of tumor cells [25,26]. Adipose tissue (abdominal fat, in the first place) produces nearly 30% of IL-6 present in the systemic circulation [27]. Our study intended to clarify whether excess bodyweight has any influence on IL-6 serum level in patients with gastrointestinal neoplasms.

Materials and Methods

Between 2014 and 2016, we studied 245 patients undergoing surgery for gastrointestinal (esophageal, hepatic, gallbladder, gastric, pancreatic, large bowel, rectal, and small intestinal) tumors at the 1st Department of Surgery of Semmelweis University (Budapest, Hungary).

The exclusion criteria were as follows: age under 18 years, inflammatory disorders (including pneumonia, wound infection, cholecystitis, peripheral cannula sepsis, endocarditis, urinary tract infection, Crohn's disease, ulcerative colitis), thromboembolic events (e.g. deep vein thrombosis, pulmonary embolism, myocardial infarction), steroid therapy, anemia (hemoglobin level <120 g/L), and artificial nutrition. Demographical and clinical data were accumulated. Preoperative blood samples were obtained closest to the time of surgery, for the measurement of IL-6 levels at the Institute for Laboratory Medicine of Semmelweis University, using an ADVIA 2120 hematology analyzer.

The same blood sample was used for the measurement of white blood cell count, and of hemoglobin concentration. BMI values were determined preoperatively, with a calibrated calculator.

Statistical Analysis

During the statistical analysis of study data, correlation was calculated using Spearman, Wilcoxon, and Pearson correlation

***Corresponding author:** Zsolt Baranyai, 1st Department of Surgery, Semmelweis University Budapest, Hungary, Tel: +3630/4500-388; E-mail: barazso@gmail.com

Received February 27, 2019; **Accepted** March 19, 2019; **Published** March 26, 2019

Citation: Jósa V, Brodszky V, Zrubka Z, Zaránd A, Mezei T, et al. (2019) The Relationship between IL-6 and Nutritional Status in Patients with Gastrointestinal Tumors. *J Oncol Res Treat* 4: 138.

Copyright: © 2019 Jósa V, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation:

coefficients. Means were compared with the paired t-test. Statistical analyses were performed at a 95% confidence level.

Based on the foregoing, 118 patients altogether had been excluded in the first place because of the presence of various inflammatory processes and thus, 127 patients were evaluated. Our study cohort was heterogeneous with regard to the types and stages of the variety of gastrointestinal tumors (Table 1). A relatively large proportion of these patients had a colorectal or a pancreatic tumor.

Among the study parameters, the extreme elevation of IL-6 levels was observed only in patients with gallbladder tumors. However, mean white blood cell count was also above the upper limit of the normal range; moreover, it is not possible to draw any meaningful conclusion from only two cases. The proportions of lean/normal-weight and of overweight/obese patients studied were similar. We did not find significant differences among the individual BMI categories, regarding neither the study parameters, nor the BMI categories by body-weight (Table 2).

The univariate analysis showed a significant correlation of IL-6 level with white blood cell count and hemoglobin level, but not between BMI and IL-6 level. The multivariate analysis confirmed a similar relationship: neither hemoglobin concentration, nor WBC influence the correlation between serum IL-6 levels and BMI (Table 3).

Discussion

Oncologists, researchers, and epidemiologists have demonstrated a relationship between excess body weight and the risk of malignancy. Hyperglycemia of 10- to 20-year duration is associated with 44-percent increase, whereas preexisting, elevated, abnormal fasting blood glucose level is accompanied by a 57-percent increase of cancer risk [28]. Moreover, the malignant tumors occurring in obese patients have

a worse prognosis [29]. The pathological background of the relationship inc-9.6

Citation:

In the microenvironment of the tumor, IL-6 supports tumorigenesis by direct action on the modulation of the intrinsic and extrinsic activity of tumor cells [44]. Basically, tumor cells produce IL-6 to promote their own survival and dissemination, whereas they do not depend on the paracrine release of IL-6 produced by stromal cells. In tumor cells, the possible stimulatory effect of IL-6 is mediated by the activation of a number of signaling pathways. IL-6 stimulates the proliferation and survival of tumor cells by activating the Ras/Raf/MEK/MAPK, PI3K/AKT, and JAK/STAT pathways through the phosphorylation of gp130 tyrosine residues [22,45]. The majority of the genes involved in the regulation of cell survival and proliferation, including Bcl-2, Bcl-xL, Mcl-1, Fas, cyclin D1, cyclin E1, and p21 are direct targets of STAT-3 along with pro-proliferation transcription factors (such as c-Myc, c-Jun, and c-FOS). In tumor cells, STAT-3 activation is mediated by the autocrine production and the paracrine secretion of IL-6 by stromal and immigrant inflammatory cells [13-16,46,47]. IL-6 has been found to mediate multiple hemopoietic effects by shifting stem cells from the G₀ to the G₁ stage of the cell cycle, and thereby to induce their proliferation and to enhance their responsiveness to additional hematopoietic factors. The latter include IL-3, IL-4, G-CSF, M-CSF, and GM-CSF [48]. The autocrine production of IL-6 by non-stem cells activates the JAK1/STAT-3 signaling pathway, which plays an important role in the transformation of non-stem cells to stem cell-like cells by up-regulating Oct-4 (a stem cell marker) [49]. Thus, IL-6 not only induces tumor cell proliferation, but also maintains their population, and this induces tumor recurrence.

C

According to our findings, the level of circulating IL-6 was not dependent on BMI in patients with various gastrointestinal tumors. It is suggested that in these cases, IL-6 production by adipose tissue is negligible as compared with cytokine expression by the tumor and its environment. Similar to chronic disorders and to long-standing inflammation, tumor-associated anemia greatly elevates IL-6 level [50,51]. Refractory anemia can develop as a consequence of pre-existing protein-energy loss [52]. A dramatic increase of IL-6 level has been found in sepsis [39]. In view of the foregoing, we also checked the hemoglobin concentrations and the white blood cell counts of our patients; however, neither of these parameters influenced the correlation between BMI and IL-6 [53].

Our method for measuring BMI has its limitations. BMI informs about the patient's height-related body mass; however, it tells nothing about the type of the tissue accrual underlying the latter [54]. Therefore, while it is a reliable indicator of obesity at population level, body mass index is of limited value when it is determined in individuals. The measurement of circulating leptin or adiponectin levels, or the assessment of body fat percentage (e.g. by computed tomography) would be much more appropriate for evaluating obesity [55]. Notwithstanding this, we believe that our findings are of potential interest, and motivate us for further research in this subject.

References

1. Renehan AG, Soerjomataram I, Tyson M, Egger M, Zwahlen M, et al. (2010) Incident cancer burden attributable to excess body mass index in 30 European countries. *Int J Cancer* 126: 692-702.
2. Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, et al. (2014) Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5.24 million UK adults. *Lancet* 384: 755-765.
3. Duan XF, Tang P, Shang XB, Jiang HJ, Zhao Q, et al. (2017) High Body Mass Index Worsens Survival in Patients with Esophageal Squamous Cell Carcinoma after Esophagectomy. *Dig Surg* 34: 319-327.
4. Cleary MP, Grossmann ME (2009) Minireview: Obesity and breast cancer: the estrogen connection. *Endocrinology* 150: 2537-2542.
5. Engin A (2017) Obesity-associated Breast Cancer: Analysis of risk factors. *Adv Exp Med Biol* 960: 571-606.
6. Michaud DS (2016) Obesity and Pancreatic Cancer. *Recent Results Cancer Res* 208: 95-105.
7. Pischon T, Nimsch K (2016) Obesity and Risk of Cancer: An Introductory Overview. *Recent Results Cancer Res* 208: 1-15.
8. Connor EV, Raker CA, Clark MA, Stuckey AR (2017) Obesity risk awareness in women with endometrial cancer. *Arch Gynecol Obstet* 295: 965-969.
9. Renehan AG, Tyson M, Egger M, Heller RF, Zwahlen M (2008) Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet* 371: 569-578.
10. De Vita F, Romano C, Orditura M, Galizia G, Martinelli E, et al. (2001) Interleukin-6 serum level correlates with survival in advanced gastrointestinal cancer patients but is not an independent prognostic indicator. *J Interferon Cytokine Res* 21: 45-52.
11. Waldner MJ, Foersch S, Neurath MF (2012) Interleukin-6-a key regulator of colorectal cancer development. *Int J Biol Sci* 8: 1248-1253.
12. Miura T, Mitsunaga S, Ikeda M, Shimizu S, Ohno I, et al. (2015) Characterization of patients with advanced pancreatic cancer and high serum interleukin-6 levels. *Pancreas* 44: 756-763.
13. Bode JG, Nimmesgern A, Schmitz J, Schaper F, Schmitt M, et al. (1999) LPS and TNFalpha induce SOCS3 mRNA and inhibit IL-6-induced activation of STAT3 in macrophages. *FEBS Lett* 463: 365-370.
14. Nagasaki T, Hara M, Nakanishi H, Takahashi H, Sato M, et al. (2014) Interleukin-6 released by colon cancer-associated fibroblasts is critical for tumour angiogenesis: anti-interleukin-6 receptor antibody suppressed angiogenesis and inhibited tumour-stroma interaction. *Br J Cancer* 110: 469-478.
15. Nolen BM, Marks JR, Ta'san S, Rand A, Luong TM, et al. (2008) Serum biomarker profiles and response to neoadjuvant chemotherapy for locally advanced breast cancer. *Breast Cancer Res* 10: R45.
16. Oh K, Lee OY, Shon SY, Nam O, Ryu PM, et al. (2013) A mutual activation loop between breast cancer cells and myeloid-derived suppressor cells facilitates spontaneous metastasis through IL-6 trans-signaling in a murine model. *Breast Cancer Res* 15: R79.
17. Lin MT, Juan CY, Chang KJ, Chen WJ, Kuo ML (2001) IL-6 inhibits apoptosis and retains oxidative DNA lesions in human gastric cancer AGS cells through up-regulation of anti-apoptotic gene mcl-1. *Carcinogenesis* 22: 1947-1953.
18. Jourdan M, Veyrune JL, De Vos J, Redal N, Couderc G (2003) A major role for Mcl-1 antiapoptotic protein in the IL-6-induced survival of human myeloma cells. *Oncogene* 22: 2950-2959.
19. Chang CH, Hsiao CF, Yeh YM, Chang GC, Tsai YH, et al. (2013) Circulating interleukin-6 level is a prognostic marker for survival in advanced nonsmall cell lung cancer patients treated with chemotherapy. *Int J Cancer* 132: 1977-1985.
20. Grivennikov S, Karin E, Terzic J, Mucida D, Yu GY, et al. (2009) IL-6 and Stat3 are required for survival of intestinal epithelial cells and development of colitis-associated cancer. *Cancer Cell* 15: 103-113.
21. Yeoh GC, Ernst M, Rose-John S, Akhurst B, Payne C, et al. (2007) Opposing

Citation: