



Figure 1





A J I

C J D

References

1. Hamilton W (2009) The CAPER studies: five case-control studies aimed at

identifying and quantifying the risk of cancer in symptomatic primary care patients. *Br J Cancer* 101: S80-S86.

2. Evans T, Sany O, Pearmain P, Ganesan R, Blann A, et al. (2011) Differential trends in the rising incidence of endometrial cancer by type: data from a UK population-based registry from 1994 to 2006. *Br J Cancer* 104: 1505-1510.

3. Abdel-Rahman M, Stockton D, Rachel B, Hakulinen T, Coleman MP, et al. (2009) What if cancer survival in Britain were the same as in Europe: how many deaths are avoidable? *Br J Cancer* 101: S115-S224.

4. Parker C, Hippiusley-Cox J, Coupland C, Vinogradova Y (2007) Rectal and postmenopausal bleeding: consultation and referral of patients with and without severe mental health problems. *Br J Gen Pract* 57: 371-376.

5. Khan NF, Harrison SE, Rose PW (2010) Validity of diagnostic coding within the General Practice Research Database: a systematic review. *Br J Gen Pract*.

6. Herrett E, Thomas SL, Schoonen WM, Smeeth L, AJ (2010) Validation and validity of diagnoses in the General Practice Research Database: a systematic review. *Br J Clin Pharmacol* 69: 4-14.

7. Hamilton W, Kernick D (2007) Clinical features of primary brain tumours: a case-control study using electronic primary care records. *Br J Gen Pract* 57: 695-699.

8. Robinson KM, Ottesen B, Christensen KB, Krasnik A (2009) Diagnostic delay experienced among gynecological cancer patients: a nationwide survey in Denmark. *Acta Obstet Gynecol Scand* 88: 685-692.