

# Tissue-Specific Differences in Mesenchymal Stromal Cells Influence Homing and Function

Christen L. Ebens<sup>1\*</sup>, Julia A. Riedl<sup>2</sup> and Jakub Tolar<sup>1,3</sup>

<sup>1</sup>*Department of Pediatrics, Division of Blood & Marrow Transplant and Cellular Therapy, University of Minnesota, Minneapolis, MN, USA*

<sup>2</sup>*Medical Scientist Training Program, University of Minnesota, Minneapolis, MN, USA*

<sup>3</sup>*Stem Cell Institute, University of Minnesota, Minneapolis, MN, USA*

\*Corresponding author:

subfamily member 5, is a surface glycoprotein that identifies (and allows for purification of) a population of DSCs with

- dystrophic epidermolysis bullosa and partial rescue via immunosuppressive dermal mesenchymal stem cells. *Lab Invest* 97: 1218-1224.
22. Beken SV, de Vries JC, Meier-Schiesser B, Meyer P, Jiang D, et al. (2019) Newly Defined ATP-binding cassette subfamily b member 5 positive dermal mesenchymal stem cells promote healing of chronic iron-overload wounds via secretion of interleukin-1 receptor antagonist. *Stem Cells* 37: 1057-1074.
23. Tappenbeck N, Schroder HM, Niebergall-Roth E, Hassinger F, Dehio U, et al. (2019) In vivo safety profile and biodistribution of GMP-manufactured human skin-derived ABCB5-positive mesenchymal stromal cells for use in clinical trials. *Cyotherapy* 21: 546-560.