" To be or Not to be "; the Choice will Always be ours, Percussions of Chronic Physical Illnesses on Pediatric Mental Health

Department of Pediatrics, Faculty of Medicine, Ain Shams University, Egypt

* Eman Ahmed Zaky, Professor of Pediatrics and Head of Child Psychiatry Clinic, Department of Pediatrics, Faculty of Medicine, Ain Shams University, Egypt, Tel: 002-01062978734; E-mail: eman2aky@hotmail.com

August 01, 2016,

August 01, 2016,

August 08, 2016

Pediatric chronic physical illnesses

Although it varies widely from one country to another, the overall prevalence of pediatric chronic physical illnesses approximately ranges from 10% to 20% [10]. On the other hand, it is estimated that 7 out of 10 c W visits to a primary care practitioner are due to chronic physical diseases [11]. YgY diseases are associated with variable degrees of impaired functions of XJ YfYbhbody organs or systems as in cases of poorly controlled bronchial asthma, chronic rheumatic heart disease, congenital heart disease, autoimmune disorders, chronic hemolytic anemias, endocrinopathies, and chronic hepatic and or renal diseases

Because of the advances in health care delivery all over the world, there is a growing concern about the mental consequences of pediatric chronic physical disorders. Chronic illnesses in childhood usually include pain, discomfort, scholastic absences, poor scholastic achievement, dependence on XJ YFVbh medications, frequent health care visits and hospitalizations, social incompetence, and poor selfesteem; all of which increase the risk of developing internalizing and externalizing mental disorders [12,13].

Percussions of pediatric chronic physical illnesses on mental health

Y disability and dysfunction that are commonly encountered with chronic illnesses were found to be attributed to the concomitant anxiety and or depression rather than the physical manifestations of such illnesses [14].

Reciprocal relationship between chronic illnesses and anxiety

It is believed that both chronic illnesses and anxiety interact with each other in ways which are likely to precipitate and or exacerbate their manifestations. On one hand, clinical anxiety increases the risk of development of medical complaints and or exaggeration of physical symptoms via either biological mediators (as hormonal or autonomic changes), psychological mechanisms (as biased symptom perception or defective coping), or a combination of both (as impaired immunity in cases of chronic worry and or excessive avoidance behaviors). On the other hand, the physical manifestations of chronic illnesses and or associated pain, dependence on medications, and or repeated hospital admissions may result in extreme fear about safety or separation from parents that eventually ends with school refusal or may lead to increased scanning and obsessions concerning physical symptoms ending with increased anxiety and panic responses to both physiological body changes and manifestations of physical diseases xixely ind dSTysical dixim

GEEAN IN BOT IN SALAR THE Calibratic Children [24] with g[b] Wahim prevalent anxiety disorders [25].

hidtt len itdas" Medi

Impact of rheumatic disorders and arthritis on pediatric mental health

Pediatric rheumatologic disorders are usually severer than the same disorders of adult onset and might lead to g[b] Wbh psychiatric morbidity with increased risk for internalizing problems especially anxiety and depression. Such high risk for psychiatric morbidity can be attributed to an autoimmune process, underlying vasculitis, and or treatment with steroids, immune suppressive medications, and or immune modulators which are potential causes of psychiatric adversities [26,27].

On the other hand, arthritis is usually associated with restriction of physical activities and might even end with physical disability that is considered as a high risk factor for the development of anxiety and or depression [20,28].

Corticosteroid therapy

Corticosteroids are indicated in a wide variety of pediatric physical diseases. Yin have been demonstrated to have g[b] Wibh neuropsychiatric side Y Wig as cognitive disorders, behavioral changes, depression, and even psychosis. Practitioners should warm caregivers about these adversities before subscribing steroids and other similar drugs to children with X] Yfybhchronic illnesses [29].

What is resilience?

Resilience is XY bYX as "the power or ability to return to the original form, position, etc....

- Fitzgerald M, Beggs S (2001) Y neurobiology of pain: developmental aspects. Neuroscientist 7: 246-257.
- 13 Jackson PL, Vessey JA, Schapiro N (2004) Primary care of the child with a chronic condition. St Lois, Mosby pp 184-191.
- 14. Kessler RC, Ormel J, Demler O, Stang PE (2003) Comorbid mental disorders account for the role impairment of commonly occurring chronic physical disorders Results from the national comorbidity survey. Journal of Occupational and Environmental Medicine 45: 1257-1266
- 15 Sareen J. Campbell DW, Leslie WD, Malisza KL, Stein MB, et al. (2007) Striatal function in generalized social phobia: a functional magnetic resonance imaging study. Biol Psychiatry 61: 396-404.
- 16 Chavira DA, Garland AF, Daley S, Hough R (2008) Yimpact of medical comorbidity on mental health and functional health outcomes among children with anxiety disorders. J Dev Behav Pediatr 29: 394-402.
- 17. Katon WJ, Richardson L, Lozano P, McCauley E (2004) Yrelationship of asthma and anxiety disorders. Psychosom Med 66: 349-355
- 18 Bloom B, Cohen RA (2007) Summary health statistics for U.S. children: National Health Interview Survey, 2006 Vital Health Stat 10 1-79.
- 19. Zaky EA, Shaheen MA, Amr NH, Said SM (2015) Psychosocial Dfc 'Y and Quality of Life in Children with Bronchial Asthma; A Comparative Study of Self Reports and Parent Proxy Reports of Peds QL[™]Inventory. Indian Journal of Applied Research 5: 33-39.
- 20 Stuber ML, Shemesh E, Saxe GN (2003) Posttraumatic stress responses in children with life-threatening illnesses. Child Adolesc Psychiatr Clin N Am 12: 195-209.
- 21. Lane DA, Millane TA, Lip GY (2013) Psychological interventions for depression in adolescent and adult congenital heart disease. Cochrane Database Syst Rev: CD004372.
- 22. Cox D, Lewis G, Stuart G, Murphy K (2002) A cross-sectional study of the prevalence of psychopathology in adults with congenital heart disease. JPsychosom Res 52:65-68
- 23 Amillategui B, Mora E, Calle JR, Giralt P (2009) Special needs of children with type 1 diabetes at primary school: perceptions from parents, children, and teachers. Pediatr Diabetes 10: 67-73

- 24 Nardi L, Zucchini S, D'Alberton F, Salardi S, Maltoni G, et al. (2008) Quality of life, psychological adjustment and metabolic control in youths with type 1 diabetes: A study with self and parent report questionnaires. Pediatric Diabetes 9: 496-503
- 25 Northam EA, Matthews LK, Anderson PJ, Cameron FJ, Werther GA (2005) Psyciatric morbidity and health outcome in type 1 diabetes-