

Upper Abdominal Surgery in Patients with Coeliac Artery Stenosis

Kyoji Ito, Nobuyuki Takemura^{*}, Ryo Oikawa, Fuyuki Inagaki, Fuminori Mihara and Norihiro Kokudo

Department of Surgery, National Centre for Global Health and Medicine, Shinjuku-ku, Tokyo, Japan

***Corresponding author:** Nobuyuki Takemura, Hepato-Biliary-Pancreatic Surgery Division, Department of Surgery, National Centre for Global Health and Medicine, 1-21-1 Toyama, Shinjuku-ku, Tokyo 162-8655, Japan, Tel +81-3-3202-7800; Fax +81-3-5273-0113; E-mail: nkokudo@hosp.ncgm.go.jp

Received date: May 11, 2020; **Accepted date:** May 26, 2020; **Published date:** June 02, 2020

Copyright: © 2020 Takemura N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits

after

difficulty

significance

effect



CA-aorta
LGA-aorta
Reconstruction using vascular graft
CHA-aorta
CA-aorta
IPDA-SPA bypass
Left iliac artery-SPA bypass

Angioplasty by IVR

Balloon dilation or stent insertion

Less invasive, anytime available in perioperative periods Difficulty in severe stenosis or complete obstruction of CA

signif cance

T e

Citation: Ito K, Takemura N, Oikawa R, Inagaki F, Mihara F, et al. (2020) Upper Abdominal Surgery in Patients with Coeliac Artery Stenosis. *Atheroscler Open Access* 5: 129.
