

Useful Method to Monitor Cerebral Infarction in Atherosclerotic Patients without Atrial Fibrillation by the Combination of Carotid Intima-Media Thickness, Cardio-Ankle Vascular Index, and Plasma D-dimer

Shigeru Hayashi^E

^ECorresponding author: Ueda Shigeru, Department of Internal Medicine, Nagoya University Graduate School of Medicine, Showa-ku, Nagoya 466-8550, Japan. E-mail: shigeru-hayashi@med.nagoya-u.ac.jp

ZLWKRXW QHXURORJLFDO V\PSWRPV ZHUH DSSURSULDWHO\ WUHDWHG GXULQJ WK

G L v H Z U H R E W D L Q H G 7 D E O H 0 R U H R Y H U 7 D E O H V K R Z V W K D W ' G L P H U
R I V X E J U R X S % J P / Z D V I R O G K L J K H U W K D Q W K D W R I
V X E J U R X S % - J ' P / Z K G L Q H U H Q F H V % 8 1 & U
} E U L Q R & H Q S O E D Q G + E Z H U H V P D O O D Q G R W K H U O D E R U D W R U \ G D W D G L G
Q R Q M v H U

| | | | |
|---------------------|---------------------|-------------|----|
| Ó:[`]íÓéíÓéáí { ^ É | Ó:[`]íÓéíÓéáí { ^ É |] | |
| ÓÉ*^Aç^Aæ*D | ÍÉÍAÀÍÈI | ÍÍHÀHÀÍÈ | ÞÙ |
| Tæ AØØ^ { æ ^ | GØFI | GÍØI Ì | ÞÙ |
| ÓTØíç*Ø { Ø | GHÉGÁÍHÉG | GHÉFÁÍHÉI | ÞÙ |
| ÙÓÚíç { P*D | FÍGÈÍÁÍGÉÉÍ | FÍLÉGÁÍFJÉÍ | ÞÙ |
| ÓÓÚíç { P*D | ÍHÉGÁÍJÉJ | ÍÍÉGÁÍJÉJ | ÞÙ |
| ÓCEØ | JÉHÀÍÈJ | JÉÍÀHÆH | ÞÙ |
| ÓTVÁç { D | FÉÍÀHÆHÈ | FÉHÌçT^åæ}D | |

IUHTXHQWO\ RFFXUUHG LQ VHW\W\HDO\LD\HCKHURVFOHURWLF SDWLHQWV

,QRXH + 1R]DZD 7 2NXPXUD . -RQJ 'DH / 6KLPL]X \$ HW DO