pathognomonic features, often presenting as an isolated, asymptomatic la enlarges. Imaging studies play a pivotal role in determining the tumor's size, relationships. The definitive diagnosis is established through histopathological Complete excision of the tumor is the sole guarantee to prevent recurrence, instances of relapse.

Undo geno a ane he ia, c  $\mathbf{m}$  e e  $\mathbf{n}$  gica  $\mathbf{e}$  e ec  $\mathbf{i}$   $\mathbf{n}$   $\mathbf{f}$ , ag . no, e / a  $\mathbf{e}$   $\mathbf{f}$   $\mathbf{e}$  med b a o a co, ica inci i  $\mathbf{n}$ . Ana mica and ah gica e a mina i  $\mathbf{n}$  c  $\mathbf{n}$   $\mathbf{e}$  med he diagn  $\mathbf{i}$   $\mathbf{f}$ , ag . no, e . ch/ ann ma.

## Discussion

Schrann mai a a e ne e genic benign me de e ed a he e en e f no e heah, and ab . ne hi d f .chrann ma ccein. he head and nec e egi n, ag . no e .chrann mai an e e e me e a e me [1]. e diagn i i di c ., i i cinica and adi gica, i e cai n i n . f l . a e n i n.


e cinica e e e nai ni n. ah gn nic. M. en, i i an i a ed, a naic, ao co ica na. ha grad a ino ea e in i e.

e i e f hi ma ma ead. ha ngea c me e i nand n n-eci c ign c cha ha ngea di c m f . e e n d n hagia.

When co, ica ma i a ciaed / ih d . h nia fi.iao a ca c · d · a a . i, ag . no, e · h/ ann ma i m · e i e · be ca · ed [2]. I maging · a · an i m · · · an · e in de o mining · he · i e and · a · ha ngea ca i n f · m · · ·

Diagnai i made ana mah gica na gica ecimen, in he ee ence f Schrann ce. f di een meh g and gani ai n, Imm n hi chemi. h / ed minan S100 e eine e e i n [3,4].

C me e e cii n f.he me i he n /a . e e en ec e ence. C n e ai n f.he e igina no e i en . .. ib e, f /ed b e e habitai n e ed ce c m icai n , /ih an e ce en ence.



**Figure 1:** Axial section of a cervico-thoracic CT scan with injection of contrast product showing a hypodense left laterocervical mass which is slightly enhanced after injection of contrast product.

**Figure 2:** Coronal reconstruction of a cervico-thoracic CT scan showing a hypodense left laterotracheal mass that displaces the common carotid artery but remains permeable.

\*Corresponding author: Sara Ez-zaky, Central Radiology Department, Ibn Sina Hospital, Morocco, E-mail: sara92ezzaky@gmail.com

Received: 02-Jan-2024, Manuscript No: roa-24-125072, Editor assigned: 05-Jan-2024, Pre-QC No: roa-24-125072 (PQ), Reviewed: 19-Jan-2024, QC No: roa-24-125072, Revised: 26-Jan-2024, Manuscript No: roa-24-125072 (R), Published: 31-Jan-2024, DOI: 10.4172/2167-7964.1000526

Citation: Ez-zaky S, Imrani K, Jellal S, Nassar I, Billah NM (2024) Vagus Nerve Schwannoma: A Rare Origin of Dysphonia and Dysphagia. OMICS J Radiol 13: 526.

Copyright: © 2024 Ez-zaky S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Final diagnosis

Vag . 📭 e. ch/ ann 📭

## Differential diagnosis list

E hagea - 🐞 •

Ade 🐧 🍃 a h

Vag . , e . a aga gg i 🚜a

## References

- Chai CK, Tang IP, Prepageran N, FRCSEdin ORL, Jayalakshmi P, et al. (2012) An Extensive Cervical Vagal Nerve Schwannoma: A Case Report. Med J Malaysia 67: 343.
- Najib Benmansour, Yasine Elfadl, Amal Bennani, Mustapha Maarouf, Leila Chbani, et al. (2013) Schwannome cervical du nerf vague: Stratégies diagnostique et thérapeutique. Pan African Medical Journal 14: 1.
- Behuria S, Rout TK, Pattanayak S (2015) Diagnosis and management of schwannomas originating from the cervical vagus nerve. Ann R Coll Surg Engl 97: 92-97.
- Kanatas A, Mücke T, Houghton D, Mitchell DA (2009) Schwannomas of the head and neck. Oncol Rev 3: 107-111.