



# Congenital Adenomatoid Cystic Malformation: Expectant Management Outcomes

Sandra Acevedo Gallegos<sup>1</sup>, María José Rodríguez Sibaja, Juan Manuel Gallardo Gaona, Jose Antonio Ramírez Calvo, Berenice Velázquez Torrez and Dulce María Camarena Cabrera

Department of Maternal-Fetal Medicine, National Institute of Perinatology, Mexico

<sup>\*</sup>Corresponding author: Sandra Acevedo Gallegos, Department of Maternal-Fetal Medicine, National Institute of Perinatology, Mexico, Tel: +525555209900; E-mail: dracevedo\_sandra@yahoo.com.mx

Received date: August 19, 2018; Accepted date: September 06, 2018; Published date: September 13, 2018

Copyright: © 2018 Gallegos SA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

**Objective:** Our purpose was to describe the evolution and perinatal outcome of Congenital Adenomatoid Cystic Malformation (CCAM) according to prenatal sonographic features in expectantly managed fetuses.

**Materials and Method:** A retrospective review of all prenatally diagnosed CCAM cases from 1995 to 2017 at a perinatal referral center in Mexico City was conducted.

**Results:** Twenty-seven cases were identified. Mean gestational age at the time of diagnosis was 26.5 weeks. According to Stocker's classification, there were eight type 1 (29.6%), ten type 2 (30%) and nine type 3 (33.4%) cases. No bilateral lesions were found. Hydrops was observed in two cases (7.4%). Lesion dimensions remained stable throughout gestation in most cases (63%), ultrasonographic regression was observed in nine fetuses (33.3%). There were no elective pregnancy terminations. Fetal demise was reported in two cases. Mean gestational age at birth was 37.1 weeks. Most newborns required only usual neonatal care. After birth, two cases showed progressive deterioration which lead to early neonatal death.

**Conclusion:** CCAM is a relatively rare abnormality of lung development, which can be accurately diagnosed by prenatal ultrasound. Conservative management is appropriate in most cases, with a favorable outcome and a chance of regression of up to one-third of the cases.

## Keywords:

Congenital Adenomatoid Cystic Malformation

classification

This

software

Text

finding

Figure

Text

identified,

Text

Text

five

Author's address

Maternal age (years) <sup>a</sup>	26 (± 6.0)
Parity <sup>b</sup>	1 (1-8)
Gestational age at the time of diagnosis <sup>a</sup>	26.5 (± 4.1)
Gestational age at birth <sup>a</sup>	37.1 (± 3.1)

Birth weight, g<sup>a</sup>

2,817 (± 598)



dif er

frst-line

Efect

Ter

Ter

af er