

Canine Distemper in Dog-Symptoms and Treatment

Department of Biomedical Sciences, University of Gondar, Ethiopia

Defersha T, Department of Biomedical Sciences, University of Gondar, Ethiopia, E-mail: defershat@vetmed.edu.com

Defersha T (2021) Canine Distemper in Dogs-Symptoms and Treatment. J Vet Med Health 6: e108

September 03, 2021;

September 17, 2021;

September 24, 2021

© 2021 Defersha T, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Introduction

Distemper is a highly contagious viral disease of dogs. It is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies. The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies.

The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies. The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies.

The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies.

1. Fever and cough
2. Nasal discharge and diarrhea
3. Lethargy and anorexia

The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies. The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies.

The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies.

Symptoms

The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies. The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies.

Stage 1

The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies. The disease is characterized by a wide range of clinical signs, including fever, cough, nasal discharge, and diarrhea. The disease is caused by the canine distemper virus (CDV) and is highly contagious. It is a major cause of mortality in dogs, especially in puppies.