



Letter

Chronic drug use is the point at which you utilize lawful or illicit substances in manners you shouldn't. You could take more than the customary portion of pills or use another person's remedy. You might manhandle medications to feel better, ease pressure, or keep away from the real world. It is accounted for that liquor incited mind issues can frequently be revised with legitimate treatment. Forbearance from liquor for months or years can help somewhat x thinking capacities,

2. Tharayil BS, Gangadhar BN, Hiremani RM, Thirthalli J, (2008) Double-blind randomized controlled study comparing short-term efficacy of bifrontal and bitemporal electroconvulsive therapy in acute mania. *Bipolar Disord* 10:701-707.
3. Dawson DA, Goldstein RB, Patricia Chou S, June Ruan W, Grant BF (2008) Age at first drink and the first incidence of adult-onset DSM-IV alcohol use disorders. *Alcohol Clin Exp Res* 32: 2149-2160.
4. The UK ECT Review Group (2003) Efficacy and safety of electroconvulsive therapy in depressive disorders: a systematic review and meta-analysis. *The Lancet* 361:799-808.
5. Shieh TY, Chang YY, Lee CY, Lu B, (2016) Patterns of betel quid, cigarette, and alcohol use, and their correlates with betel quid cessation in a male inmate population. *Substance Use Misuse* 51:471-478.