

1 2 3 4 5 6 7 8 9 10 11

In the ca age, the bifacial is more common in female than in male. This is due to the fact that the biological change in accommodation. In the end of the ing di ance e i emen, a k e med, and a m leng h [8,9]. The bifacial cec in l d lead an e en gea e di a i among men and w men in e m f nced e bifacia e cec ed e bifacia [8].

1 2 3 4 5 6 7 8 9 10 11

The bifacial is a facie e. If e c n ide he nced e facie e in he ca e f e bifacia, indi ec l i a ec e bifacia ign & m m. L a e f nced e h e ia can ccc ea need f e bifacia cec in bef e he age f 40.

1 2 3 4 5 6 7 8 9 10 11

The bifacial is a facie e. If e c n ide he nced e facie e in he ca e f e bifacia, indi ec l i a ec e bifacia ign & m m. L a e f nced e h e ia can ccc ea need f e bifacia cec in bef e he age f 40.

1 2 3 4 5 6 7 8 9 10 11

The bifacial is a facie e. If e c n ide he nced e facie e in he ca e f e bifacia, indi ec l i a ec e bifacia ign & m m. L a e f nced e h e ia can ccc ea need f e bifacia cec in bef e he age f 40.

In the f m f bifacial cec in n d egmen, D- egmen, e ec i e, ibbn egmen, and c ed egmen de ign a e a ailable.

Rnd egmen k k bifacial i he chea e bifacial i h egmen f 22 f 38mm. I i ed f all e f eeing in di ance and nea bjec. A he ical cen e f he e gla e lie a he ge me ical cen e f he egmen, i gi e an e a ba e-d n i m e ec. I gi e an e ce eld f i e. Image j m i m e in hi e (Fig e 1).

Fla bifacial i al called a D- egmen. Segmen heigh a ie f m 22 f 45 mm m l ed i 28mm. A ide eading a ea gi e a be e e ec in nea i n b i a ec di ance i n m all di ance egmen. Image j m i neg i able ince he ical cen e i nea he di ance egmen. C ed egmen imila a egmen i h

Progressive additional lenses are prescribed by the optometrist for the elderly (Elderly). These lenses are designed to be available for reading and distance vision. Distance vision has a wide reading area and distance area. Progressive lenses are in the intermediate zone and have a wide reading area. Distance lenses are in the distance zone and have a wide reading area.

Conclusion

Presbyopia is a global problem, affecting billions of people worldwide in aging populations. Corrective lenses are available for all the people who are affected by this condition. However, the use of glasses is not always preferred because of the cost and the inconvenience of wearing them. The use of contact lenses is also not always preferred because of the cost and the inconvenience of wearing them. The use of contact lenses is also not always preferred because of the cost and the inconvenience of wearing them.

References

1. Holden BA, Fricke TR, Ho SM (1999) Global Vision Impairment Due to Uncorrected Presbyopia. Arch Ophthalmol 126: 1731-1739.
2. Kamali A, Whitworth JA, Ruberantwari A, Mulwany F, Acakara M, et al. (1999) Causes and prevalence of non-vision impairing ocular conditions among a rural adult population in SW Uganda. Ophthalmic Epidemiol 6: 41-8.
3. Wharton KR, Ylton RL (1986) Visual characteristics of rural Central and South Americans. J Am Optom Assoc 57: 426-30.
4. Burke AG, Patel I, Munoz B, Kayongoya A, McHiwa W (2006) Population-based study of presbyopia in rural Tanzania. Invest Ophthalmol Vis Sci 47: 1005-1010.