

-
1. Glucosylsphingosine is a key biomarker of Gaucher disease. *Am J Hematol* 11: 1082-1089.
 2. Bultron G, Kacena K, Pearson D, Boxer M, Yang M, et al. (2010) The risk of Parkinson's disease in type 1 Gaucher disease. *J Inherit Metab Dis* 33: 167-173.
 3. The human glucocerebrosidase gene and pseudogene: structure and evolution. *Genomics* 4: 87-96.
 4. locus on chromosome 1q21: implications for Gaucher disease. *Genome Res* 7: 1020-1026.
 5. Grabowski GA (2012) Gaucher disease and other storage disorders. *Hematology Am Soc Hematol Educ Program* 2012: 13-8.
 6. Koprivica V, Stone DL, Park JK, Callahan M, Frisch A, et al. (2000) Analysis Gaucher disease. *Am J Hum Genet* 66: 1777-1786.
 7. Mohebbi N, Wagner CA (2018) Pathophysiology, diagnosis and treatment of inherited distal renal tubular acidosis. *J Nephrol*, 31: 511-522.
 8. Clinical and molecular aspects of distal renal tubular acidosis in children. *Pediatr Nephrol* 32: 987-996.
 9. Prenatal Diagnosis of Tay-Sachs Disease. *Methods Mol Biol* 1885: 233-250.
 10. Acid stimulation of the citrate transporter NaDC-1 requires Pyk2 and ERK1/2 signaling pathways. *J Am Soc Nephrol* 29: 1720-1730.
-