

Keywords: Concoction; Sero-reconversion; HIV Patient; Herbal therapy

Introduction

Herbal remedies contain active ingredients which are parts of plants or plant materials, or combinations used to treat a multitude of ailments throughout the world. It was estimated that most Africans use herbal remedies [1]. The severity of illnesses, availability and low cost prompted many to seek for alternative therapies outside orthodox medicines. This explains why some HIV infected patients seek for herbal therapies due to its un-curable nature.

It was estimated in 2010 that 63% of the HIV infected people in the world live in sub-Saharan Africa [2]. Since the discovery of acquired immunodeficiency syndrome (AIDS) linked with HIV as the causative agent in early 1980s, Nigeria still battles with the scourge of this dreadful infection [3,4]. It was estimated that there are over 3 million HIV infected people in Nigeria thus the second country in the world with largest population of people infected with HIV infection [2].

The use of herbal remedies for terminal illness is very common and the fact that HIV infection had no cure prompted many to seek for traditional medicines and spiritual solutions. People infected with HIV infection in Nigeria started looking for super-natural solutions since 1987 when the first case was reported [5]. Although free highly active antiretroviral therapy (HAART) is widely available, many HIV patients in Nigeria are desperately looking for quicker solution to the existing problems of long duration therapy by patronising herbal therapist. Campaigns by media houses and discouragement by medical practitioners had not stopped the herbal therapists from flourishing in HIV infection treatment business.

Many herbal remedies have been documented to be useful in HIV infection. Like highly active antiretroviral therapy, many herbal remedies that have been found to inhibit one or more steps in HIV replication [6,7]. A carbohydrate derivative, pentosan poly-sulphate

had been documented to inhibit HIV tat regulatory protein (p14) that strongly activates transcription of proviral DNA [8]. *Acacia* from tropical liana plant inhibit reverse transcriptase and HIV induced cell fusion [9]. Coumarins (coumarins) from tropical forest tree (*Cassia*) was considered as non-nucleoside reverse transcriptase inhibitor in potency [10,11].

women and prostitutes especially when on distant journey? spanning through the Northern, Eastern and Western parts of Nigeria. He had blood transfusion (commercial donor) at the beginning of the illness in a private hospital. The examination revealed conscious but very weak patient with marked weight loss (evidenced by prominent zygomatic process, clavicular bones and body mass index of 17.8), skin rashes, moderate pallor, and oral thrush on the buccal cavity that was widely spread on tongue and extending to the throat. There was no significant palpable peripheral lymphadenopathy. The laboratory investigation revealed positive reaction to HIV 1 antigens (ELISA) which was confirmed with western blot as positive (P24 and gp 41 antigens). Chest X-ray, electrolytes, renal function tests, liver function tests and full blood count were not significantly impaired. However, CD4 count showed 178 cells/mm³ and viral load could not be done because PCR was not available for routine HIV diagnosis in Nigeria.

Because of the non-availability of free anti-viral agent in 2005 at the nearest teaching hospital, the patient opted for herbal remedy thus discharged himself from hospital against medical advice. The patient was encouraged to visit hospital on weekly basis for supportive medical assistance (opportunistic infections) but shown up a year a month on herbal therapy requesting for repeat HIV screening. HIV (EIA) was done and positive although, all the presented clinical features associated with HIV infection had disappeared. The investigation revealed that he had commenced a 4-month herbal concoction therapy (10ml oral paste diluted with warm water three times daily). The repeated CD4 count a year a month therapy revealed 160 cells/mm³ despite full recovery of the patient from all the presenting symptoms and signs of HIV/AIDS.

The subsequent 2nd and 3rd month HIV screening were positive with 190 and 300 cells/mm³ respectively. The fourth month (134th day on herbal therapy) CD4 count was 420 cells/mm³ but ELISA (Genscreen) HIV screening was negative. Western blot (Newlab blot 1& 2) was repeated a month thereafter and the sero-negativity was confirmed.

The sero-reversion led to review of the initial patient's diagnosis with the blood sample safely stored in teaching hospital laboratory. Several repeat diagnostic reviews (clinical and HIV testing) at different laboratories with different kits (Genscreen, Determine, Stakpack, Unigold, Newlab blot 1&2, etc) of the patient confirmed HIV/AIDS at WHO grade staging 3 (minimum). However, viral load could not be done on the stored sample. The patient was followed up quarterly with repeat CD4 count, HIV screening and confirmation for 18 months then, yearly till 2012. The patient had normal CD4 count and sero-negativity

throughout the follow up. The viral (HIV-RNA) load was commenced in 2009 and had been persistently undetectable (< 50 copies/ml) in the body fluids (blood and semen) (Table 1).

Comments

The sero-positivity of HIV patient is based on the antibody produced to HIV antigen [16]. It is assumed that when HIV antibody is detected in an individual, it is very rare to become undetectable or sero-negative irrespective of the extent of treatment. Therefore, any case of sero-reversion attracts multiple reviews on diagnosis. All the diagnostic criteria, history and examination were reviewed and found to be accurate for the diagnosis of HIV infection at WHO stage 4 for the above reported case. What is responsible for full recovery and sero-reversion of HIV infection within 5 months of herbal therapy remained a million's question. The above case and associated question would have been answered if the patient was sero-reverted spontaneously before commencing herbal therapy [17].

Since the discovery of HIV infection in 1980's, the common fact is that HIV/AIDS is not curable but could be managed leading patient to long life therapy. Despite the fact that human immunodeficiency virus disappeared in blood circulation (aviraemia) after 3-4 months of intensive therapy with highly active anti-retroviral therapy (HAART), sero-reconversion had been a very rare case. The few documented cases of sero-reconversion in HIV patient on highly antiretroviral therapy (HAART) were at early stage and acute phase of infection [18,19]. Spontaneous sero-reversion did not occur in this patient unlike earlier study [20]. Contrary to the expectation, the above patient who was at terminal phase of HIV infection recovered and became sero-reverted which was not associated with death [21]. The sustained EIA and western blot sero-negativity in this patient had been a major concern because partial sero-reversion had been documented [19,22]. The sustained 6 years period of sero-reversion of this HIV patient is in contrary to temporary sero-negativity in earlier documented case [23].

The explanation for sustained complete sero-reversion in the above case could only be attributed to herbal therapy unlike immunosuppressive

References

1. WHO (2002) Traditional Medicine Growing Needs and Potential. WHO Policy Perspectives on Medicines, World Health Organization, Geneva 1-6.
2. UNAIDS/WHO (2010) The Millenium Goals report 2010.
3. Barré-Sinoussi F, Chermann JC, Rey F, Nugeyre MT, Chamaret S, et al. (1983) Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). *Science* 220: 868-871.
4. Gallo RC, Sarin PS, Gelmann EP, Robert-Guroff M, Richardson E, et al. (1983) Isolation of human T- cell leukemia virus in acquired immune deficiency syndrome (AIDS). *Science* 220: 865-867.
5. Abalaka JO (2004) Attempts to cure and prevent HIV/AIDS in central Nigeria between 1997 and 2002: opening a way to a vaccine-based solution to the problem? *Vaccine* 22: 3819-3828.
6. De Clercq E (2000) Current lead natural products for the chemotherapy of human immunodeficiency virus (HIV) infection. *Med Res Rev* 20: 323-349.
7. Kong JM, Goh NK, Chia LS, Chia TF (2003) Recent advances in traditional plant drugs and orchids. *Acta Pharmacol Sin* 24: 7-21.
8. Matthée G, Wright AD, König GM (1999) HIV reverse transcriptase inhibitors of natural origin. *Planta Med* 65: 493-506.
9. Watson K, Gooderham NJ, Davies DS, Edwards RJ (1999) Interaction of the transactivating protein HIV-1 tat with sulphated polysaccharides. *Biochem Pharmacol* 57: 775-783.
10. Dharmaratne HR, Tan GT, Marasinghe GP, Pezzuto JM (2002) Inhibition of HIV-1 reverse transcriptase and HIV-1 replication by *Calophyllum* coumarins and xanthenes. *Planta Med* 68: 86-87.
11. Yu D, Suzuki M, Xie L, Morris-Natschke SL, Lee KH (2003) Recent progress in the development of coumarin derivatives as potent anti-HIV agents. *Med Res Rev* 23: 322-345.
12. Lu W, Wen R, Guan C, Wang Y, Shao J, et al. (1995) A report on 8 seronegative converted HIV/AIDS patients with traditional Chinese medicine. *Zhongguo Zhong xi yi jie he za zhi Zhongguo Zhongxiyi jiehe zazhi Chinese journal of*