

## Abstract

The Present case was an Immune reconstitution inflammatory syndrome (IRIS) affecting the central nervous system (CNS) in patient with HIV and tuberculosis in HIV patient initiated on antiretroviral therapy. This case identified as the patient was visited Life Care Clinic, at Aurangabad. The patient was newly initiated on antiretroviral therapy (ART), The IRIS was diagnosed based on National AIDS Control Organization guidelines. A typical Mycobacterial CNS-IRIS in AIDS patient was represented, the increase in CD4 count in a rapid way and biopsy showing on encephalitis our diagnosis of IRIS was confirmed. The patient was started on prednisolone and his hemiparesis

**Keywords:** Immune reconstitution inflammatory syndrome (IRIS); Central nervous system (CNS) CD4/TB/HIV/ART

## Introduction

Patients with HIV/Tuberculosis (TB) co-infection who are on anti-TB treatment and highly active antiretroviral therapy (HAART) will develop an exacerbation of symptoms, signs or radiological manifestations of TB that are not due to recurrence or relapse of their tuberculosis [1-4]. The IRIS is defined as the occurrence or manifestation of new opportunistic infections or existing opportunistic infections within six weeks to six months after initiating the antiretroviral therapy with increase in the CD4 cells. The IRIS is a complex and a not yet completely well-understood phenomenon that is seen most commonly in HIV infection, the aetiology of these immune reconstitution inflammatory syndrome (IRIS) reactions is unknown, but it is presumed that they occur, as a consequence of HAART-related reconstitution of immunity and leading to an abnormal immune response to antigens released by dead or dying bacilli. Mycobacteria account for approximately 40% of all cases of infective IRIS in patients

biopsy showing on encephalitis our diagnosis of IRIS was confirmed.

The patient was started on prednisolone and his hemiparesis improved mass effect decreased and patient went home walking then after six months and the Patient weighed about 55kgs and his CD4 count was 359 cells/mm

<sup>3</sup> and is asymptomatic this confirms Immune reconstitution inflammatory syndrome (IRIS) affecting the central nervous system (CNS) in patient with HIV and tuberculosis.

---

**\*Corresponding author:** Dr. Shivaji Kashinath Jadhav, Ph.D, Scientist, Dr.Prabhakar Kore, Basic Science Research Centre, VK Institute of Dental Sciences, Campus KLE University, Nehru nagar, Belgaum-590010, Karnataka, India, Tel: +91 9740341860; E-mail: [shiv\\_jk@yahoo.co.in](mailto:shiv_jk@yahoo.co.in)

**Received** February 28, 2013; **Published** March 26, 2013

**Citation:** Udaykumar Gopinath J, Shivaji Kashinath J, Manohar Gangappa P (2013) Immune Reconstitution Inflammatory Syndrome (IRIS) Affecting the Central Nervous System (CNS) In Patient with HIV and Tuberculosis on Antiretroviral Therapy. 2: 653 doi:[10.4172/scientificreports.653](https://doi.org/10.4172/scientificreports.653)

**Copyright:** © 2013 Udaykumar Gopinath J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

---

12. Kumarasamy N, Chaguturu S, Mayer KH, Solomon S, Yepthomi HT, et al. (2004) Incidence of immune reconstitution syndrome in HIV/tuberculosis co infected patients after initiation of generic antiretroviral therapy in India. *J Acquir Immune Defc Syndr* 37: 1574-1576.
13. Narita M, Ashkin D, Hollender ES, Pitchenik AE (1998) Paradoxical worsening of tuberculosis following antiretroviral therapy in patients with AIDS. *Am J Respir Crit Care Med* 158: 157-161.
14. Mínguez C, Roca B, González-Miño C, Simón E (2000) Superior vena cava syndrome during the treatment of pulmonary tuberculosis in an HIV-1 infected patient. *J Infect* 40: 187-189.
15. Furrer H, Malinverni R (1999) Systemic inflammatory reaction after starting highly active antiretroviral therapy in AIDS patients treated for extrapulmonary tuberculosis. *Am J Med* 106: 371-372.
16. Fishman JE, Saraf-Lavi E, Narita M, Hollender ES, Ramsinghani R, et al. (2000) Pulmonary tuberculosis in AIDS patients: transient chest radiographic worsening after initiation of antiretroviral therapy. *AJR Am J Roentgenol* 174: 43-49.
17. Hejazi N, Hassler W (1997) Multiple intracranial tuberculomas with atypical response to tuberculostatic chemotherapy: literature review and a case report. *Acta Neurochir (Wien)* 139: 194-202.
18. Cheng VC, Ho PL, Lee RA, Chan KS, Chan KK, et al. (2002) Clinical spectrum of paradoxical deterioration during antituberculosis therapy in non-HIV-infected patients. *Eur J Clin Microbiol Infect Dis* 21: 803-809.
19. Teoh R, Humphries MJ, O'Mahony G (1987) Symptomatic intracranial tuberculoma developing during treatment of tuberculosis: a report of 10 patients and review of the literature. *Q J Med* 63: 449-460.
20. Crump JA, Tyrer MJ, Lloyd-Owen SJ, Han LY (1998) Military tuberculosis with paradoxical expansion of intracranial tuberculomas complicating human immunodeficiency virus infection in a patient receiving highly active antiretroviral therapy. *Clin Infect Dis* 26: 1008-1009.
21. Vidal JE, Cimerman S, Schiavon Nogueira R, Bonasser Filho F, Sztajnbock J, et al. (2003) Paradoxical reaction during treatment of tuberculous brain abscess in a patient with AIDS. *Rev Inst Med Trop Sao Paulo* 45: 177-178.