during the complementary feeding period. e complementary foods are o en of low nutritional quality and given in insu cient amounts. When given too early or too frequently, they displace breast milk [5]. In Ethiopia poor feeding practices and shortfall in food intake are the most important direct factors responsible for malnutrition and illness amongst children [5] and a combination of nutritionally inferior diets and improper feeding practices are major contributing factors to the development of childhood malnutrition [12]. Complementary feeding

unre ned cereals, grains, legumes, and oil seeds are a potent inhibitor of iron, zinc, and calcium absorption. Hence, to ensure absorption

Ba

It is one of the most important staple food crops in the high lands of Ethiopia. Currently, barley is widely consumed as a food grain with desirable nutritional contents consumed as a foods and snacks are increasingly available, driven by recent research ndings, which show that barley bers contains beta–glucans and tocotrineols, chemical agents known to lower serum cholesterol level [31]. In Ethiopia barley is the h most important cereal crop next to maize, tef, sorghum and wheat. It is the staple food grain for Ethiopian high lands who mange the crop with indigenous technologies and utilized di erent parts of the plants for various purposes.

Barley is used as our, as semolina, and as whole-dehulled grain.

Citation: Temesgen M (2013)

phytate degradation can a ect the nutritional value of high phytate diet. Furthermore, enzymatic hydrolysis of phytate in the gastrointestinal tract leads to the formation of speci c isomer of inositol phosphates formed during food processing or degradation in the gastrointestinal

Citation: Temesgen M (2013)