

Open Access Scientific Reports

Case Report

Open Access

Keywords: Breastfeeding; Baby-friendly; Exclusive breastfeeding; Health care

Introduction

Lactation is a natural process that allows the mother to provide her infant with the best source of nutrition and protection. The World Health Organization (WHO) and the United Nations Children's Fund (UNICEF) recommend exclusive breastfeeding for the first 6 months of life. In 2010, the global prevalence of exclusive breastfeeding was 44.7%. In the United States, the prevalence of exclusive breastfeeding at 6 months was 16.3%. The World Health Organization (WHO) estimates that 11,300 children die each day from preventable causes related to inadequate breastfeeding. The World Health Organization (WHO) estimates that 11,300 children die each day from preventable causes related to inadequate breastfeeding. The World Health Organization (WHO) estimates that 11,300 children die each day from preventable causes related to inadequate breastfeeding.

The Baby-Friendly Initiative (BFHI) was launched in 1991 to improve breastfeeding practices. The Baby-Friendly Initiative (BFHI) was launched in 1991 to improve breastfeeding practices. The Baby-Friendly Initiative (BFHI) was launched in 1991 to improve breastfeeding practices.

adequate breastfeeding practices are essential for the health and survival of infants. The World Health Organization (WHO) estimates that 11,300 children die each day from preventable causes related to inadequate breastfeeding. The World Health Organization (WHO) estimates that 11,300 children die each day from preventable causes related to inadequate breastfeeding.

Citation:

Citation: Toma TS, Rea MF (2012)

... a.) ... ab ... e ... d ... f ... e ... a ...
 ... e ... e ... a ... ca ... be ... e ... WHO a ... UNICEF
 ... e ... d ... a ... e ... a ... f ... e ... a ... bec ... a BFH,
 ... a ... a ... d ... f ... e ... e ... a ... d ... e ... be ... e ... a ... a ... ed ... e ... Bra ... a ...
 ... M ... f ... Hea ... ad ... ed ... a ... a ... a ... d ... e ... e ... f ... a ...
 ... c ... e ... ed ... be ... a ... ed ... ce ... a ... e ... a ... f ... f ... f ...
 ... e ... e ... f ... e ... e ... a ... a ... d ... a ... e ... f ... be ... ed ... be
 ... e ... e ... a ... ed ... b ... e ... e ... a ... e ... a ... a ... I ... be ... a ... f ...
 ... f ... e ... e ... a ... e ... e ... e ... e ... a ... f ... ab ... a ... d ... e ... A ... a ...
 ... f ... e ... ec ... e ... da ... a ... a ... e ... a ... e ... c ... f ... c ... e ...
 ... f ... a ... d ... e ... a ... de ... a ... e ... ec ... e ... ed ... f ...
 ... a ... de ... a ... a ... f ... e ... BFHI ... e ... ab ... e ... a ... e ... b ... e ... d ... c ...
 ... a ... d ... e ... f ... e ... a ... e ... a ... e ... a ... e ...

References

1. United Nations Children's Fund (UNICEF) (2008) World's Children 2009 - Maternal and Newborn Health. Brasilia: 158.
2. Edmond KM, Zandoh C, Quigley MA, Amenga-Etego S, Owusu-Agyei S, et al. (2006) Delayed breastfeeding initiation increases risk of neonatal mortality. *Pediatrics* 117: 380-386.
3. Labbok M (2007) Breastfeeding and Baby-Friendly Hospital Initiative: more important and with more evidence than ever. *J Pediatr (Rio J)* 83: 99-101.
4. World Health Organization (WHO) (2001) Scientific evidence of the ten steps to successful breast feeding. Brasilia: Pan American Health Organization.
5. Kramer MS, Chalmers B, Hodnett ED, Sevkovskaya Z, Dzikovich I, et al. (2001) The Probit Study Group. Promotion of Breastfeeding Intervention Trial. *Trials* 2: 11.