



(F, 1). . . . . D  
620<sup>1</sup> 2 . . . . . G  
07 08 12 35 59 15 E.  
68,000<sup>1</sup> 2, . . . . .  
610<sup>1</sup> 2 . . . . . F . . . . . (F) . . . . .  
450 . . . . .  
190<sup>1</sup>

...; A ... D ...  
... (F, 2. B, C, D ... F).  
... (F, 2. A).  
... *A. hydrophila*.  
...  
... *Aeromonads* ... H,  
...  
*Aeromonas* ... 28 C ...  
... 38-41 C.  
... 24.6 C. 27.2 C  
... D ...  
14 ...  
... *Aeromonas*  
... 27% ... A ... B ...  
*A. veronii* biovar *sobria* ... 6 ... 11  
... B ... *Aeromonas hydrophila*  
... E ... 15 ...  
... *A. hydrophila* ...



## References

1. Conte FS (2004) Stress and the welfare of cultured fish. *Appl Anim Behav Sci* 86: 205-223.
2. Plumb JA (1994) Health Maintenance and Principal Microbial Diseases of Cultured Fishes. *Animal diseases*, 254.
3. González-Serrano CJ, Santos JA, García-López ML, Otero A (2002) Virulence markers in *Aeromonas hydrophila* and *Aeromonas veronii biovar sobria* isolates from freshwater fish and from a diarrhoea case. *J Appl Microbiol* 93: 414-419.
4. Doukas, Athanassopoulou, Karagouni, Dotsika (1998) Short communication *Aeromonas hydrophila* infection in cultured sea bass, *Dicentrarchus labrax* L., and