


& G G F D U P G % J M V U J P O P O U I F Q) B O E 5 J U S B U
. F E J D J O F T

Viviane Santos da Silva Pierro ¹, Lucianne Cople Maia ^{1*}, Thiago Isidro Vieira ¹, Monique Santana Candreva ¹, Michele Villardi ² and Lucio Mendes Cabral ²

¹Department of Pediatric Dentistry and Orthodontics, School of Dentistry, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

²Department of Medicaments, Pharmacy School, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil



Keywords: Tooth erosion; Pharmaceutical preparations; Hydrogen-Materials and Methods

ion concentration; Titrimetry; Child

Introduction

Dental erosion has been demanding increasing attention as a result of the increasing prevalence of dental caries reduction in many societies [1]. It can be defined "as chronic loss of dental hard tissue that is chemically etched away from the tooth surface by acid and/or chelation without bacterial involvement" [2,3]. It is assumed that the main etiological factors are acids of intrinsic

medicines analyzed.

The pediatric syrup medicines chosen for this study (Dimetapp®, Wyeth, Sao Paulo, Brazil and Claritin®, Schering-Plough, Vila Olímpia, Brazil) were selected based on a previous study [12], which has pointed out these medicines as presenting the worst results with regard to pH and titratable acidity taken together. Control solutions of citric acid with similar baseline pH values to those of the medicines tested were also evaluated.

The pH measurements throughout the study were made using a pH electrode connected to a digital pH-meter (Quimis Q-400HM, Diadema, Brazil). The electrode was calibrated at the start of each session using standard buffers of pH 4.1 and 6.86. Five milliliters of the newly opened medicines and controls, which were at room temperature, were placed in a beaker and stirred using a non-heating magnetic stirrer until a stable reading was obtained. All measurements were obtained in triplicate and three bottles from each pediatric syrup medicine presenting different serial numbers were analyzed.

Further five milliliters from each bottle of syrup medicine and both control solutions of citric acid were diluted with water (pH 6.48 ± 0.12) in a proportion of 1:0.5 and 1:1, and pH of these dilutions were measured as described earlier.

The titratable acidity was determined in triplicate by using the same pH-meter. To detect the end point of each medicine and control (both in undiluted and diluted forms), five milliliters of each substance were titrated with 0.05 M sodium hydroxide (NaOH) solution, using

[10]. From measurements of hardness and by SEM and pro lo metric analysis of bovine teeth and tests of the dissolution of hydroxyapatite *in vitro*, another study [24] verified that citric acid was more erosive than malic acid, being a potent erosive agent because of its ability to chelate calcium in hydroxyapatite, thus increasing enamel's rate of dissolution on exposure to the acid. In the present study, both control solutions were citric acid with similar baseline pH values to those of the medicines tested. The acid's choice was based on the medicines' labels, which denoted the presence of citric acid in the medicines' formulation. Our results confirmed the ability of citric acid to resist pH changes because of controls' high values of titratable acidity. When compared to the medicines tested, control solutions of citric acid presented titratable acidic values three to fourfold greater than medicines' values.

Other factors related with the ingested substances themselves could also modify erosion patterns, such as the general chemical composition of the solutions, which may modify the degree of enamel dissolution [25]. Ions like calcium, phosphate and fluoride have a protective effect against erosion [23]. Therefore, previous studies have proposed the reduction of the erosive potential of beverages by modifying the amount and type of acid used in beverage formulations and/or supplementing with calcium and phosphate [14-16,24].

Nevertheless, product re-formulation can be difficult to be proposed because additives could promote further effects on other ingredients in the drinks [14]. Therefore, an alternative to products' modification could be products' dilution as tested by previous researchers [17] in a study that aimed to examine the effect of dilution on the potential erosive properties of some diluting drinks. Their results showed that only titratable acidity fell considerably as the drink was progressively diluted, and little effect was observed on pH values. However, these results were based on high dilution rates (concentrations ranged from undiluted to one part drink in 100,000 parts water), which were not applicable to the usual consumption of these drinks. On the other hand, our findings demonstrated that dilution influenced neither pH nor titratable acidity, but it is important to point out that our study intended to verify the effect of dilution on the erosive potential of highly acidic pediatric syrup medicines and, therefore, the proposed dilution rates could not be more than twice the volume prescribed (which corresponds to a dilution rate of 1:1) because of the probable diet of children in drinking medicines' volumes greater than these.

The etiology of dental erosion can also be influenced by biological and behavioral factors. According to a previous study [23], "unusual eating, drinking and swallowing habits, for example holding an acid beverage in the mouth before swallowing, increase the contact time of an acid substance with the teeth and thus increase the risk of erosion" [23]. It is also true for acidic syrup medicines when used with a high frequency of ingestion and consumption at bedtime, considering that oral clearance is compromised during sleep [8-9,13]. In this context, both pediatric syrup medicines tested in this study could increase the risk of tooth erosion for their consumers. Besides being highly acidic, Claritin® is an antihistamine and one of its side effects is the reduction of salivary flow rates, which impairs the protective effect of saliva against erosion. With regard to Dimetapp®, it presented pH values and titratable acidic values even worse than Claritin's and its label information recommends its use up to 6 times per day.

Eroded enamel is more susceptible to wear by tooth-brushing and to toothpaste abrasion [26,27]. Therefore, it is not advisable to brush the teeth immediately after consuming acidic foods or drinks because it could result in tooth abrasion. Instead, a delay in tooth-brushing would allow the remineralizing action of saliva on the eroded enamel [27].

Conclusion

Within the limitations of this study, the dilution rates tested did not improve the pH and titratable acidity of the acidic pediatric syrup medicines analyzed. Therefore, as dilution did not reduce the erosive properties of syrup medicines tested considering our results, recommendation on immediate water rinse and delayed tooth-brushing after syrup medicines' ingestion could be proposed at the time of prescription. Further studies evaluating alternative preventive strategies for dental erosion due to these highly acidic syrup medicines should be conducted.

Acknowledgement

The authors also would like to thank CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico - Conselho Nacional de Desenvolvimento Científico e Tecnológico - Fundação Carlos Chagas de Amparo a Pesquisa do Estado do Rio de Janeiro) for the financial support.

References

- Lussi A, Jaeggi T, Zero D (2004) The role of diet in the aetiology of dental erosion. *Caries Res* 38: 34-44.
- PIHO G, HGWDO HURVLRQ, HQLWLRQ FOD... *Sci* 104: 151-155.
- ten Cate JM, Imfeld T (1996) Dental erosion, summary. *Eur J Oral Sci* 104: 241-244.
- Lussi A, Jaeggi T, Schaffner M (2002) Diet and dental erosion. *Nutrition* 18: 780-781.
- Dugmore CR, Rock WP (2004) A multifactorial analysis of factors associated with dental erosion. *Br Dent J* 196: 283-286.
- Moss SJ (1998) Dental erosion. *Int Dent J* 48: 529-539.
- Nunn JH, Ng SK, Sharkey I, Coulthard M (2001) The dental implications of chronic use of acidic medicines in medically compromised children. *Pharm World Sci* 23: 118-119.
- Pierro VS, Abdelnur JP, Maia LC, Trugo LC (2005) Free sugar concentration and pH of paediatric medicines in Brazil. *Community Dent Health* 22: 180-183.
- Costa CC, Almeida IC, Costa Filho LC (2006) Erosive effect of an antihistamine-FRQWDLQLQJ V\UXS RQ SULPDU\ HQDPHO DQG LWV Paediatr Dent 16: 174-180.
- Maguire A, Baqir W, Nunn JH (2007) Are sugars-free medicines more erosive than sugars-containing medicines? An *in vitro* study of paediatric medicines with prolonged oral clearance used regularly and long-term by children. *Int J Paediatr Dent* 17: 231-238.

21. Grenby TH, Phillips A, Desai T, Mistry M (1989) Laboratory studies of the dental properties of soft drinks. *Br J Nutr* 62: 451-464.
22. Larsen MJ, Nyvad B (1999) Enamel erosion by some soft drinks and orange juices relative to their pH, buffering effect and contents of calcium phosphate. *Caries Res* 33: 81-87.
- 23.